

**From Quantum Chaos
To
Anderson Localization**

Boris Altshuler

Columbia University

NEC Laboratories America

RANDOM MATRICES

$N \times N$ matrices with random matrix elements. $N \rightarrow \infty$

Dyson Ensembles

Matrix elements

real

complex

2×2 matrices

Ensemble

orthogonal

unitary

symplectic

RANDOM MATRIX THEORY

E_α

- spectrum (set of eigenvalues)

$$\delta_1 \equiv \langle E_{\alpha+1} - E_\alpha \rangle$$

- mean level spacing

RANDOM MATRIX THEORY

E_α

- spectrum (set of eigenvalues)

$$\delta_1 \equiv \langle E_{\alpha+1} - E_\alpha \rangle$$

- mean level spacing

$\langle \dots \rangle$

- ensemble averaging

RANDOM MATRIX THEORY

$$E_\alpha$$

- spectrum (set of eigenvalues)

$$\delta_1 \equiv \langle E_{\alpha+1} - E_\alpha \rangle$$

- mean level spacing

$$\langle \dots \rangle$$

- ensemble averaging

$$s \equiv \frac{E_{\alpha+1} - E_\alpha}{\delta_1}$$

- spacing between consecutive eigenvalues

RANDOM MATRIX THEORY

$$E_\alpha$$

- spectrum (set of eigenvalues)

$$\delta_1 \equiv \langle E_{\alpha+1} - E_\alpha \rangle$$

- mean level spacing

$$\langle \dots \rangle$$

- ensemble averaging

$$s \equiv \frac{E_{\alpha+1} - E_\alpha}{\delta_1}$$

- spacing between consecutive eigenvalues

$$P(s)$$

- distribution function

RANDOM MATRIX THEORY

$$E_\alpha$$

$$\delta_1 \equiv \langle E_{\alpha+1} - E_\alpha \rangle$$

$$\langle \dots \rangle$$

$$s \equiv \frac{E_{\alpha+1} - E_\alpha}{\delta_1}$$

$$P(s)$$

- spectrum (set of eigenvalues)

- mean level spacing

- ensemble averaging

- spacing between consecutive eigenvalues

- distribution function

Spectral Rigidity
Level repulsion

$$P(s = 0) = 0$$

$$P(s \ll 1) \propto s^\beta \quad \beta=1,2,4$$

Noncrossing rule (theorem) $P(s=0) = 0$

Suggested by Hund (*Hund F. 1927 Phys. v.40, p.742*)

Justified by von Neumann & Wigner (*v. Neumann J. & Wigner E. 1929 Phys. Zeit. v.30, p.467*)

Usually textbooks present a simplified version of the justification due to Teller (*Teller E., 1937 J. Phys. Chem 41 109*).

Arnold V. I., 1972 Funct. Anal. Appl.v. 6, p.94

*Mathematical Methods of Classical Mechanics
(Springer-Verlag: New York), Appendix 10, 1989*


***Arnold V.I., Mathematical Methods of Classical Mechanics
(Springer-Verlag: New York), Appendix 10, 1989***

In general, a multiple spectrum in typical families of quadratic forms is observed only for two or more parameters, while **in one-parameter families of general form the spectrum is simple for all values of the parameter. Under a change of parameter in the typical one-parameter family the eigenvalues can approach closely, but when they are sufficiently close, it is as if they begin to repel one another. The eigenvalues again diverge, disappointing the person who hoped, by changing the parameter to achieve a multiple spectrum.**

$$\hat{H}(x) \Rightarrow E_{\alpha}(x)$$

*Arnold V.I., Mathematical Methods of Classical Mechanics
(Springer-Verlag: New York), Appendix 10, 1989*

In general, a multiple spectrum in typical families of quadratic forms is observed only for two or more parameters, while in one-parameter families of general form the spectrum is simple for all values of the parameter. Under a change of parameter in the typical one-parameter family the eigenvalues can approach closely, but when they are sufficiently close, it is as if they begin to repel one another. The eigenvalues again diverge, disappointing the person who hoped, by changing the parameter to achieve a multiple spectrum.



Reason for $P(s) \rightarrow 0$ when $s \rightarrow 0$:

$$\hat{H} = \begin{pmatrix} H_{11} & H_{12} \\ H_{12}^* & H_{22} \end{pmatrix}$$

$$E_2 - E_1 = \sqrt{(H_{22} - H_{11})^2 + |H_{12}|^2}$$

small

small

small

1. The assumption is that the matrix elements are statistically independent. Therefore probability of two levels to be degenerate vanishes.
2. If H_{12} is real (orthogonal ensemble), then for s to be small two statistically independent variables $((H_{22} - H_{11})$ and $H_{12})$ should be small and thus $P(s) \propto s \quad \beta = 1$

Reason for $P(s) \rightarrow 0$ when $s \rightarrow 0$:

$$\hat{H} = \begin{pmatrix} H_{11} & H_{12} \\ H_{12}^* & H_{22} \end{pmatrix}$$


$$E_2 - E_1 = \sqrt{(H_{22} - H_{11})^2 + |H_{12}|^2}$$

small

small

small

1. The assumption is that the matrix elements are statistically independent. Therefore probability of two levels to be degenerate vanishes.
2. If H_{12} is **real (orthogonal ensemble)**, then for s to be small **two statistically independent** variables ($(H_{22} - H_{11})$ and H_{12}) should be small and thus $P(s) \propto s$ $\beta = 1$
3. **Complex H_{12} (unitary ensemble)** \implies both $Re(H_{12})$ and $Im(H_{12})$ are statistically independent \implies **three** independent random variables should be small $\implies P(s) \propto s^2$ $\beta = 2$




**Gaussian
Orthogonal
Ensemble**

Orthogonal
 $\beta=1$

Unitary
 $\beta=2$

Symplectic
 $\beta=4$

Poisson – completely uncorrelated levels



$N \times N$ matrices with random matrix elements. $N \rightarrow \infty$

Spectral Rigidity
Level repulsion

$$P(s \ll 1) \propto s^\beta \quad \beta = 1, 2, 4$$

Dyson Ensembles

Realizations

<u>Matrix elements</u>	<u>Ensemble</u>	β	
real	orthogonal	1	T-inv potential
complex	unitary	2	broken T-invariance (e.g., by magnetic field)
2×2 matrices	symplectic	4	T-inv, but with spin-orbital coupling

Finite size quantum physical systems

Atoms

Nuclei

Molecules

.

.

.




Quantum
Dots

ATOMS

Main goal is to classify the eigenstates in terms of the quantum numbers

NUCLEI

For the nuclear excitations this program does not work



N. Bohr, Nature
137 (1936) 344.

ATOMS


Main goal is to classify the eigenstates in terms of the quantum numbers

NUCLEI


For the nuclear excitations this program does not work

E.P. Wigner
(*Ann.Math*, v.62, 1955)

Study spectral **statistics** of a **particular** quantum system
- a given nucleus




N. Bohr, Nature
137 (1936) 344.



Particular
nucleus

^{166}Er



Spectra of
several
nuclei
combined
(after
spacing)
rescaling
by the
mean level

ATOMS

Main goal is to classify the eigenstates in terms of the quantum numbers

NUCLEI

For the nuclear excitations this program does not work

E.P. Wigner
(*Ann.Math*, v.62, 1955)

Study spectral **statistics** of a **particular** quantum system - a given nucleus

Random Matrices	Atomic Nuclei
<ul style="list-style-type: none">• <i>Ensemble</i>• <i>Ensemble averaging</i>	<ul style="list-style-type: none">• <i>Particular quantum system</i>• <i>Spectral averaging (over α)</i>

Nevertheless

Statistics of the nuclear spectra are almost exactly the same as the Random Matrix Statistics

T-invariance (CP) violation - **crossover**
between Orthogonal and Unitary
ensembles

Q

■
■

Why the random matrix theory (RMT) works so well for nuclear spectra

?

Q ■ ■ *Why the random matrix theory (RMT) works so well for nuclear spectra*



Original answer:

*These are systems with a **large number of degrees of freedom**, and therefore the “complexity” is high*

Later it became clear that

there exist very “simple” systems with as many as 2 degrees of freedom ($d=2$), which demonstrate RMT - like spectral statistics

Classical Dynamical Systems with d degrees of freedom

Integrable Systems

The variables can be separated and the problem reduces to d one-dimensional problems



d integrals of motion

Classical ($\hbar = 0$) Dynamical Systems with d degrees of freedom

Integrable Systems

The variables can be separated and the problem reduces to d one-dimensional problems




d integrals of motion

Examples

1. A ball inside rectangular billiard; $d=2$

- **Vertical** motion can be separated from the **horizontal** one

- **Vertical** and **horizontal** components of the momentum, are both integrals of motion



Classical ($\hbar = 0$) Dynamical Systems with d degrees of freedom

Integrable Systems

The variables can be separated and the problem reduces to d one-dimensional problems




d integrals of motion

Examples

1. A ball inside rectangular billiard; $d=2$

- **Vertical** motion can be separated from the **horizontal** one


- **Vertical** and **horizontal** components of the momentum, are both integrals of motion



2. Circular billiard; $d=2$

- **Radial** motion can be separated from the **angular** one

- **Angular** momentum and **energy** are the integrals of motion



Classical Dynamical Systems with d degrees of freedom

Integrable Systems

The variables can be separated \Rightarrow d one-dimensional problems \Rightarrow d integrals of motion

Rectangular and circular billiard, Kepler problem, . . . ,
1d Hubbard model and other exactly solvable models, . .

Classical Dynamical Systems with d degrees of freedom

Integrable Systems

The variables can be separated \Rightarrow d one-dimensional problems \Rightarrow d integrals of motion

Rectangular and circular billiard, Kepler problem, . . . ,
1d Hubbard model and other exactly solvable models, . .

Chaotic Systems

The variables **can not** be separated \Rightarrow there is only one integral of motion - energy

Examples

Classical Dynamical Systems with d degrees of freedom

Integrable Systems


The variables can be separated \Rightarrow d one-dimensional problems \Rightarrow d integrals of motion

Rectangular and circular billiard, Kepler problem, . . . ,
1d Hubbard model and other exactly solvable models, . .

Chaotic Systems

The variables **can not** be separated \Rightarrow there is only one integral of motion - energy

Examples



Classical Dynamical Systems with d degrees of freedom

Integrable Systems


The variables can be separated \Rightarrow d one-dimensional problems \Rightarrow d integrals of motion

Rectangular and circular billiard, Kepler problem, . . . ,
1d Hubbard model and other exactly solvable models, . .

Chaotic Systems

The variables **can not** be separated \Rightarrow there is only one integral of motion - energy

Examples



Stadium

Classical Dynamical Systems with d degrees of freedom

Integrable Systems


The variables can be separated \Rightarrow d one-dimensional problems \Rightarrow d integrals of motion

Rectangular and circular billiard, Kepler problem, . . . , 1d Hubbard model and other exactly solvable models, . .

Chaotic Systems

The variables **can not** be separated \Rightarrow there is only one integral of motion - energy

Examples



Stadium

Classical Dynamical Systems with d degrees of freedom

Integrable Systems


The variables can be separated \Rightarrow d one-dimensional problems \Rightarrow d integrals of motion

Rectangular and circular billiard, Kepler problem, . . . , 1d Hubbard model and other exactly solvable models, . .


Chaotic Systems

The variables **can not** be separated \Rightarrow there is only one integral of motion - energy

Examples



Sinai billiard



Stadium

Classical Dynamical Systems with d degrees of freedom

Integrable Systems


The variables can be separated \Rightarrow d one-dimensional problems \Rightarrow d integrals of motion

Rectangular and circular billiard, Kepler problem, . . . , 1d Hubbard model and other exactly solvable models, . .


Chaotic Systems

The variables **can not** be separated \Rightarrow there is only one integral of motion - energy


Examples



Sinai billiard



Stadium




Kepler problem
in magnetic field


Chaotic Systems

The variables **can not** be separated \Rightarrow there is only one integral of motion - energy


Examples



Sinai billiard



Stadium



Kepler problem in magnetic field



Yakov Sinai



Leonid Bunimovich





Johannes Kepler

Classical Chaos

$$\hbar = 0$$

- *Nonlinearities*
- *Lyapunov exponents*
- *Exponential dependence on the original conditions*
- *Ergodicity*



Quantum description of any System with a finite number of the degrees of freedom is a linear problem - Shrodinger equation

Q: What does it mean Quantum Chaos ?

$\hbar \neq 0$

Bohigas – Giannoni – Schmit conjecture

VOLUME 52

2 JANUARY 1984

NUMBER 1

Characterization of Chaotic Quantum Spectra and Universality of Level Fluctuation Laws

O. Bohigas, M. J. Giannoni, and C. Schmit

Division de Physique Théorique, Institut de Physique Nucléaire, F-91406 Orsay Cedex, France

(Received 2 August 1983)

It is found that the level fluctuations of the quantum Sinai's billiard are consistent with the predictions of the Gaussian orthogonal ensemble of random matrices. This reinforces the belief that level fluctuation laws are universal.

In
summary, the question at issue is to prove or dis-
prove the following conjecture: Spectra of time-
reversal-invariant systems whose classical an-
alogs are K systems show the same fluctuation
properties as predicted by GOE

$\hbar \neq 0$

Bohigas – Giannoni – Schmit conjecture

VOLUME 52

2 JANUARY 1984

NUMBER 1

Characterization of Chaotic Quantum Spectra and Universality of Level Fluctuation Laws

O. Bohigas, M. J. Giannoni, and C. Schmit

Division de Physique Théorique, Institut de Physique Nucléaire, F-91406 Orsay Cedex, France

(Received 2 August 1983)

It is found that the level fluctuations of the quantum Sinai's billiard are consistent with the predictions of the Gaussian orthogonal ensemble of random matrices. This reinforces the belief that level fluctuation laws are universal.

In
summary, the question at issue is to prove or dis-
prove the following conjecture: Spectra of time-
reversal-invariant systems whose classical an-
alogs are K systems show the same fluctuation
properties as predicted by GOE

$\hbar \neq 0$

Bohigas – Giannoni – Schmit conjecture

VOLUME 52

2 JANUARY 1984

NUMBER 1

Characterization of Chaotic Quantum Spectra and Universality of Level Fluctuation Laws

O. Bohigas, M. J. Giannoni, and C. Schmit

Division de Physique Théorique, Institut de Physique Nucléaire, F-91406 Orsay Cedex, France


(Received 2 August 1983)

It is found that the level fluctuations of the quantum Sinai's billiard are consistent with the predictions of the Gaussian orthogonal ensemble of random matrices. This reinforces the belief that level fluctuation laws are universal.

Chaotic
classical analog




Wigner- Dyson
spectral statistics



No quantum
numbers except
energy

In

summary, the question at issue is to prove or disprove the following conjecture: Spectra of time-reversal-invariant systems whose classical analogs are K systems show the same fluctuation properties as predicted by GOE



Q: What does it mean Quantum Chaos ?

Two possible definitions


Chaotic
classical
analog

Wigner -
Dyson-like
spectrum

Classical


Quantum

Integrable




Poisson

Chaotic




Wigner-Dyson



Poisson to Wigner-Dyson crossover

Important example: quantum particle subject to a random potential - disordered conductor

✱ *Scattering centers, e.g., impurities*




Poisson to Wigner-Dyson crossover

Important example: quantum particle subject to a **random potential** - disordered conductor

✧ *Scattering centers, e.g., impurities*

- As well as in the case of Random Matrices (RM) there is a luxury of ensemble averaging.
- The problem is much richer than RM theory
- There is still a lot of universality.



Anderson localization (1956)


At strong enough disorder all eigenstates are **localized** in space

Correlations due to Localization in Quantum Eigenfunctions of Disordered Microwave Cavities

Prabhakar Pradhan and S. Sridhar

Department of Physics, Northeastern University, Boston, Massachusetts 02115


(Received 28 February 2000)

***Anderson Insulator******Anderson Metal***

Poisson to Wigner-Dyson crossover

Important example: quantum particle subject to a random potential - disordered conductor

✧ *Scattering centers, e.g., impurities*




Models of disorder:
Randomly located impurities

$$U(\vec{r}) = \sum_i u(\vec{r} - \vec{r}_i)$$

Poisson to Wigner-Dyson crossover

Important example: quantum particle subject to a random potential - disordered conductor

✧ Scattering centers, e.g., impurities



Models of disorder:

Randomly located impurities

$$U(\vec{r}) = \sum_i u(\vec{r} - \vec{r}_i)$$

White noise potential


$$u(\vec{r}) \rightarrow \lambda \delta(\vec{r}) \quad \lambda \rightarrow 0 \quad c_{im} \rightarrow \infty$$

Anderson model - tight-binding model with onsite disorder

Lifshits model - tight-binding model with offdiagonal disorder

-
-
-

Anderson Model



- *Lattice - tight binding model*
- *Onsite energies ϵ_i - **random***
- *Hopping matrix elements I_{ij}*

$$-W < \epsilon_i < W$$

uniformly distributed

$$I_{ij} = \begin{cases} I & \mathbf{i} \text{ and } \mathbf{j} \text{ are nearest neighbors} \\ 0 & \text{otherwise} \end{cases}$$

Anderson Transition

$$I < I_c$$

Insulator

*All eigenstates are **localized***
Localization length ξ


$$I > I_c$$

Metal

*There appear states **extended***
all over the whole system


Localization of single-electron wave-functions:

$$\left[-\frac{\nabla^2}{2m} + U(\mathbf{r}) - \epsilon_F \right] \psi_\alpha(\mathbf{r}) = \xi_\alpha \psi_\alpha(\mathbf{r})$$



Localization of single-electron wave-functions:

$$\left[-\frac{\nabla^2}{2m} + U(\mathbf{r}) - \epsilon_F \right] \psi_\alpha(\mathbf{r}) = \xi_\alpha \psi_\alpha(\mathbf{r})$$



$d=1$; All states are localized

$d=2$; All states are localized

$d>2$; Anderson transition

Anderson Transition

$$I < I_c$$

Insulator

All eigenstates are localized
Localization length ξ

The eigenstates, which are localized at different places will not repel each other



Poisson spectral statistics

$$I > I_c$$

Metal

There appear states extended all over the whole system

Any two extended eigenstates repel each other




Wigner – Dyson spectral statistics

Zharekeshev & Kramer.

Exact diagonalization of the Anderson model


3D cube of volume 20x20x20



Quantum particle in a random potential (*Thouless, 1972*)


Energy scales

1. Mean level spacing



$$\delta_1 = 1/v_{\times} L^d$$

energy



L is the system size;

d is the number of dimensions

D is the diffusion const

2. Thouless energy

$$E_T = hD/L^2$$

E_T has a meaning of the *inverse diffusion time* of the traveling through the system or the *escape rate* (for open systems)

$$g = E_T / \delta_1$$

dimensionless
Thouless
conductance

$$g = Gh/e^2$$




Scaling theory of Localization

(Abrahams, Anderson, Licciardello and Ramakrishnan 1979)

$$g = E_T / \delta_1$$

*Dimensionless Thouless
conductance*

$$g = Gh/e^2$$



$$L = 2L = 4L = 8L \dots$$

without quantum corrections

$$E_T \propto L^{-2} \quad \delta_1 \propto L^{-d}$$


Scaling theory of Localization

(Abrahams, Anderson, Licciardello and Ramakrishnan 1979)

$$g = E_T / \delta_1$$

Dimensionless *Thouless*
conductance


$$g = Gh/e^2$$



$$L = 2L = 4L = 8L \dots$$

without quantum corrections


$$E_T \propto L^{-2} \quad \delta_1 \propto L^{-d}$$





$$\frac{d(\log g)}{d(\log L)} = \beta(g)$$

β - function

$$\frac{d \log g}{d \log L} = \beta(g)$$





Metal - insulator transition in 3D
All states are localized for $d=1,2$



Conductance g

Anderson transition in terms of pure level statistics







Correlations due to Localization in Quantum Eigenfunctions of Disordered Microwave Cavities

Prabhakar Pradhan and S. Sridhar

Department of Physics, Northeastern University, Boston, Massachusetts 02115

(Received 28 February 2000)

Integrable*Square
billiard****Chaotic****Sinai
billiard***All chaotic
systems
resemble
each other.***Disordered
localized***All integrable
systems are
integrable in
their own way***Disordered
extended*

Disordered Systems:

$$E_T > \delta_1; \quad g > 1$$

*Anderson metal;
Wigner-Dyson spectral statistics*

$$E_T < \delta_1; \quad g < 1$$

*Anderson insulator;
Poisson spectral statistics*

Q: *Is it a generic scenario for the Wigner-Dyson to Poisson crossover ?*

Speculations

Consider an *integrable* system. Each state is characterized by a *set of quantum numbers*.

It can be viewed as a point in the *space of quantum numbers*. The whole set of the states forms a *lattice* in this space.

A *perturbation* that violates the integrability provides matrix elements of the *hopping* between different sites (*Anderson model !?*)

Q: *Does Anderson localization provide a generic scenario for the Wigner-Dyson to Poisson crossover ?*

Consider an *integrable* system. Each state is characterized by a *set of quantum numbers*.

It can be viewed as a point in the *space of quantum numbers*. The whole set of the states forms a *lattice* in this space.

A *perturbation* that violates the integrability provides matrix elements of the *hopping* between different sites (*Anderson model !?*)

Weak enough hopping - Localization - Poisson
Strong hopping - transition to Wigner-Dyson

The very definition of the localization is **not invariant** - one should specify in which space the eigenstates are localized.

Level statistics **is invariant**:

Poissonian statistics

\exists basis where the eigenfunctions are localized

Wigner -Dyson statistics

\forall basis the eigenfunctions are extended

Example 1

Doped semiconductor

Low concentration
of donors




Electrons are localized on
donors \Rightarrow **Poisson**

Higher donor
concentration



Electronic states are
extended \Rightarrow **Wigner-Dyson**



Example 1

Doped semiconductor

Low concentration of donors




Electrons are localized on donors \Rightarrow Poisson

Higher donor concentration



Electronic states are extended \Rightarrow Wigner-Dyson




Example 2

Rectangular billiard

Two integrals of motion

$$p_x = \frac{\pi n}{L_x}; \quad p_y = \frac{\pi m}{L_x}$$

Lattice in the momentum space




Line (surface) of constant energy

Ideal billiard

- localization in the momentum space \Rightarrow Poisson

Deformation or smooth random potential

- delocalization in the momentum space \Rightarrow Wigner-Dyson



Diffusion and Localization in Chaotic Billiards

Fausto Borgonovi,^{1,3,4} Giulio Casati,^{2,3,5} and Baowen Li^{6,7}

¹*Dipartimento di Matematica, Università Cattolica, via Trieste 17, 25121 Brescia, Italy*

²*Università di Milano, sede di Como, Via Lucini 3, Como, Italy*

³*Istituto Nazionale di Fisica della Materia, Unità di Milano, via Celoria 16, 22100, Milano, Italy*

⁴*Istituto Nazionale di Fisica Nucleare, Sezione di Pavia, Pavia, Italy*


⁵*Istituto Nazionale di Fisica Nucleare, Sezione di Milano, Milano, Italy*

⁶*Department of Physics and Centre for Nonlinear and Complex Systems, Hong Kong Baptist University, Hong Kong*

⁷*Center for Applied Mathematics and Theoretical Physics, University of Maribor, Krekova 2, 2000 Maribor, Slovenia*

(Received 29 July 1996)

$$\varepsilon \equiv \frac{a}{R}$$



$\varepsilon > 0$ **Chaotic stadium**

$\varepsilon \rightarrow 0$ **Integrable circular billiard**

Angular momentum is
the integral of motion

$$\hbar = 0; \quad \varepsilon \ll 1$$

Diffusion in the
angular momentum
space

$$D \propto \varepsilon^{5/2}$$

Localization
and diffusion
in the angular
momentum
space

Localization and diffusion in the angular momentum space

Diffusion and Localization in Chaotic Billiards

Fausto Borgonovi,^{1,3,4} Giulio Casati,^{2,3,5} and Baowen Li^{6,7}

¹Dipartimento di Matematica, Università Cattolica, via Trieste 17, 25121 Brescia, Italy

²Università di Milano, sede di Como, Via Lucini 3, Como, Italy

³Istituto Nazionale di Fisica della Materia, Unità di Milano, via Celoria 16, 22100, Milano, Italy

⁴Istituto Nazionale di Fisica Nucleare, Sezione di Pavia, Pavia, Italy


⁵Istituto Nazionale di Fisica Nucleare, Sezione di Milano, Milano, Italy

⁶Department of Physics and Centre for Nonlinear and Complex Systems, Hong Kong Baptist University, Hong Kong

⁷Center for Applied Mathematics and Theoretical Physics, University of Maribor, Krekova 2, 2000 Maribor, Slovenia

(Received 29 July 1996)

$$\varepsilon \equiv \frac{a}{R}$$



$\varepsilon > 0$ Chaotic stadium




$\varepsilon \rightarrow 0$ Integrable circular billiard

Angular momentum is the integral of motion

$$\hbar = 0; \quad \varepsilon \ll 1$$

Diffusion in the angular momentum space

$$D \propto \varepsilon^{5/2}$$



D.Poilblanc, T.Ziman, J.Bellisard, F.Mila & G.Montambaux
Europhysics Letters, v.22, p.537, 1993

1D Hubbard Model on a periodic chain

$$H = t \sum_{i,\sigma} \left(c_{i,\sigma}^+ c_{i+1,\sigma} + c_{i+1,\sigma}^+ c_{i,\sigma} \right) + U \sum_{i,\sigma} n_{i,\sigma} n_{i,-\sigma} + V \sum_{i,\sigma,\sigma'} n_{i,\sigma} n_{i+1,\sigma'}$$

$V = 0$ Hubbard model

integrable

Onsite interaction

n. neighbours interaction

$V \neq 0$ extended Hubbard model

nonintegrable

D.Poilblanc, T.Ziman, J.Bellisard, F.Mila & G.Montambaux
Europhysics Letters, v.22, p.537, 1993

1D Hubbard Model on a periodic chain

$$H = t \sum_{i,\sigma} \left(c_{i,\sigma}^+ c_{i+1,\sigma} + c_{i+1,\sigma}^+ c_{i,\sigma} \right) + U \sum_{i,\sigma} n_{i,\sigma} n_{i,-\sigma} + V \sum_{i,\sigma,\sigma'} n_{i,\sigma} n_{i+1,\sigma'}$$

$V = 0$ Hubbard model

integrable

Onsite interaction

n. neighbors interaction

$V \neq 0$ extended Hubbard model


nonintegrable

12 sites


3 particles

Zero total spin




Total momentum $\pi/6$



D.Poilblanc, T.Ziman, J.Bellisard, F.Mila & G.Montambaux
Europhysics Letters, v.22, p.537, 1993





D.Poilblanc, T.Ziman, J.Bellisard, F.Mila & G.Montambaux
Europhysics Letters, v.22, p.537, 1993




1d t - J model



D.Poilblanc, T.Ziman, J.Bellisard, F.Mila & G.Montambaux
Europhysics Letters, v.22, p.537, 1993



Chaos in Nuclei – Delocalization?



•

•

•

•

•

• • • •

•

•



**Delocalization
in Fock space**