

Generalized Julia Sets: from Cantor Bouquet to Cantor Cheese

Stefanie Hittmeyer¹, Bernd Krauskopf², and Hinke M. Osinga³

Department of Mathematics, The University of Auckland, New Zealand

¹stefanie.hittmeyer@auckland.ac.nz, ²b.krauskopf@auckland.ac.nz, ³h.m.osinga@auckland.ac.nz

Abstract

The complex quadratic family $z \mapsto z^2 + c$ is well known to give rise to artistic images and renderings of *Julia sets* — for a given complex number c , the Julia set is the boundary of the set of points z that escape to infinity under iteration. There exists a unique *critical value* c , which is the image of the critical point $z = 0$. If iterates of c remain bounded then the Julia set is connected, typically a set of closed (Jordan) curves, and if its iterations diverge to infinity then the Julia set is a Cantor set. Here, we consider a non-analytic perturbation of the complex quadratic family and its *generalized Julia sets*, which are defined in the same way. There are now infinitely many critical values that lie on a circle. Consequently, some critical values may remain bounded under iteration, while others do not. As the result, the (generalized) Julia set can be more complicated, leading to new intriguing mathematical objects that may provide novel inspirations for art.

Introduction

One of the paradigm examples in dynamical systems theory is the complex quadratic family [4] defined as

$$z \mapsto z^2 + c,$$

for $z \in \mathbb{C}$ and parameter $c \in \mathbb{C}$. This map generates a dynamical system for any fixed c by iteration: for an arbitrary choice of initial point $z = z_0 \in \mathbb{C}$, its orbit is the infinite sequence

$$\{z_0, z_1, z_2, \dots\}, \text{ with } z_i \in \mathbb{C} \text{ and } z_{i+1} = z_i^2 + c, \text{ for all } i \in \mathbb{N}.$$

In polar coordinates, $z = r e^{i\theta}$, the complex quadratic family becomes

$$r e^{i\theta} \mapsto r^2 e^{2i\theta} + c,$$

which shows that the map involves angle doubling (due to the z^2 term). The dynamics of the complex quadratic map for a given c is primarily determined by whether and which points in the complex plane remain bounded under iteration. The boundary of this set of bounded orbits is called the Julia set. It is a special property of the complex quadratic map, called the *fundamental dichotomy*, that its Julia set is connected if, and only if, the orbit of the critical value c is bounded; otherwise, if the orbit of c goes to ∞ , the Julia set is always a Cantor set, that is, it is totally disconnected [3].

In this paper, we consider the perturbation or generalization of the complex quadratic map given by

$$f_\lambda(z) = \left(1 - \lambda + \lambda |z|^2\right) \left(\frac{z}{|z|}\right)^2 + c. \quad (1)$$

Here, the parameter $c \in \mathbb{C}$ is as in the complex quadratic map, but there is now an additional (real) parameter $\lambda \in [0, 1]$ that modifies the quadratic term. This family of maps was motivated by the map introduced in [2] with $c = 1$ in the context of higher-dimensional (wild) chaotic dynamics in a Lorenz-like vector field in \mathbb{R}^5 ; see also the geometric studies in [6, 7, 12] of this type of chaotic dynamics of the map (1). As for the

quadratic map, it is easier to understand the action of the map (1) when it is written in polar coordinates, namely, as

$$re^{\vartheta i} \mapsto (1 - \lambda + \lambda r^2) e^{2\vartheta i} + c.$$

This form shows that the map (1) is essentially like the complex quadratic map, still doubling the angle ϑ , but now the radial term is $1 - \lambda + \lambda r^2$, which differs from r^2 if, and only if, $\lambda \neq 1$. Note that the map (1) is not defined for $z = 0$, but considering $r = 0$ gives the set $J_1 := \{(1 - \lambda) e^{2\vartheta i} + c \mid \vartheta \in (-\pi, \pi)\}$ as the ‘image’ of $z = 0$; note that J_1 is a circle of radius $1 - \lambda$ centered around c . (The notation J_1 comes from the literature of non-invertible maps and was also used in our earlier work [6, 7, 8, 9, 12].) Just as c is the critical value for the complex quadratic map, now J_1 plays the role of critical values for the map (1). By this we mean that the properties of the generalized Julia set of the map (1) are determined by the fate of the orbits starting on the critical circle J_1 .

In this paper, we discuss some of the specific findings reported in [8, 9] with focus on the mathematical and artistic properties of the generalized Julia sets of (1). These are again defined as the boundary of the set of points that escape to infinity (for fixed c and λ), and they can be computed in exactly the same way as for the quadratic map, namely by coloring the complex plane according to the fate of initial conditions under iteration. For the images presented here, points in the complex plane are colored yellow if their associated orbits remain bounded; otherwise, we color them blue. More specifically, we consider a grid of 1000×1000 initial conditions $z_0 = x + yi$ in the range $-1.2 \leq x, y \leq 1.2$ and decide the color based on up to $N = 500$ iterations. If during the iteration we find that $|z_i| \geq 2.4$ then the initial condition is colored blue; on the other hand, if $|z_N| < 2.4$ then the initial condition is colored yellow. This method of distinguishing initial conditions can be varied in many ways to generate artistically pleasing images; in particular, one can define a color gradient that represents the rate at which orbits diverge to ∞ , or converge to a finite attractor (if it exists).

Generalized Julia sets with novel geometric properties

Figure 1 shows generalized Julia sets of the map (1) when $c = 0.1$ (top row) and $c = 0.28$ (bottom row). The case shown in panel (a) is representative when the orbit of any point on J_1 converges to an attracting (and hyperbolic) periodic point. Just as for the complex quadratic map, the generalized Julia set is then a set of so-called Jordan curves that separate the basins of attraction of the attracting periodic point from the points that go off to infinity; in Figure 1(a) there is an attracting fixed point and the Julia set is a single Jordan curve. Similarly, if none of the points in J_1 remain bounded then the generalized Julia set is a Cantor set, that is, totally disconnected (not shown).

New types of generalized Julia sets arise for the the ambivalent case where some points on J_1 escape to infinity and other points on J_1 remain bounded. Figure 1(b) shows an example for the particular case that the critical point $z = 0$ lies *inside* J_1 . The generalized Julia now forms a *Cantor bouquet*, which is an infinite union of arcs, connected in a single so-called *explosion point*, such that the end points of the arcs are dense. The Cantor bouquet as a Julia set has been discovered before, namely, in the complex exponential family $z \mapsto \lambda e^z$; there, it exists when $\lambda < e^{-1}$, but then the explosion point lies at infinity [1, 5, 10, 11]. Panels (c) and (d) of Figure 1 show two additional ambivalent cases. As for panel (b), the generalized Julia sets are connected, but they each have fundamentally different geometric properties. For both of these cases, the critical point $z = 0$ lies *outside* J_1 and the points on J_1 that escape to infinity form a family of curve segments. At the same time, J_1 contains infinitely many points that remain bounded. The difference is that in panel (c) the map (1) does not have a finite attractor, while in panel (d) it does. When there is no finite attractor, the generalized Julia set is a *critically connected Cantor set*, which is characterized by the property that there is a countable dense set of connection points and that it contains Jordan curves that bound open regions of points that escape to ∞ . In Figure 1(d), a finite attractor exist, so that there also exist open regions




Figure 1: Generalized Julia sets (boundary of the blue region) of the map (1). Panel (a) for $c = 0.1$ and $\lambda = 0.78$ is a Jordan curve. Panel (b) for $c = 0.1$ and $\lambda = 0.6$ is a Cantor bouquet. Panel (c) for $c = 0.28$ and $\lambda = 0.91$ is a critically connected Cantor set. Panel (d) for $c = 0.28$ and $\lambda = 0.89$ is a Cantor cheese.

of points that converge to this attractor; these are, consequently, colored yellow. We call this generalized Julia set a *Cantor cheese*.

Conclusions

The map (1) exhibits more complicated generalized Julia sets than are known for the complex quadratic map. This is due to the new and third possibility that some points on the critical circle J_1 escape to infinity, while others do not. The generalized Julia sets for this ambivalent case come in a number of different variants,

and we showed here a Cantor bouquet, a critically connected Cantor set, and a Cantor cheese. There are more cases, and it is not yet clear whether all cases have been discovered; we refer to [8, 9] for more details. From an aesthetic point of view, these new types of generalized Julia sets offer opportunities for creating novel and intriguing artistic renderings. While they are just as straightforward to generate, generalized Julia sets are fundamentally different from well-known images of Julia sets for complex quadratic maps and other complex analytic maps. Hence, the map presented here can be seen as a new playground that may inspire the mathematical arts community.

References

- [1] J. M. Aarts, and L. G. Oversteegen. “The geometry of Julia sets.” *Transactions of the American Mathematical Society*, vol. 338, no. 2, 1993, pp. 897–918. <http://www.ams.org/journals/tran/1993-338-02/S0002-9947-1993-1182980-3/S0002-9947-1993-1182980-3.pdf>.
- [2] R. Bamón, J. Kiwi and J. Rivera-Letelier, “Wild Lorenz-like attractors.” arXiv 0508045, 2006. <http://arxiv.org/abs/math/0508045>.
- [3] P. Blanchard. “Complex analytic dynamics on the Riemann sphere.” *Bulletin of the American Mathematical Society*, vol. 11, no. 1, 1984, pp. 85–141. <https://www.ams.org/journals/bull/1984-11-01/S0273-0979-1984-15240-6/S0273-0979-1984-15240-6.pdf>.
- [4] R. L. Devaney. *An introduction to chaotic dynamical systems*. Westview Press, Boulder, CO, 2003.
- [5] R. L. Devaney and M. Krych. “Dynamics of $\exp(z)$.” *Ergodic Theory & Dynamical Systems*, vol. 4, no. 1, 1984, pp. 35–52. <https://doi.org/10.1017/S014338570000225X>.
- [6] S. Hittmeyer, B. Krauskopf and H. M. Osinga. “Interacting global invariant sets in a planar map model of wild chaos.” *SIAM Journal on Applied Dynamical Systems*, vol. 12, no. 3, 2013, pp. 1280–1329. <https://doi.org/10.1137/120902860>
- [7] S. Hittmeyer, B. Krauskopf and H. M. Osinga. “From wild Lorenz-like to wild Rovella-like dynamics.” *Dynamical Systems*, vol. 30, no. 4, 2015, pp. 525–542. <https://doi.org/10.1080/14689367.2015.1081677>.
- [8] S. Hittmeyer, B. Krauskopf and H. M. Osinga. “Interactions of the Julia set with critical and (un)stable sets in an angle-doubling map on $\mathbb{C}\setminus\{0\}$.” *International Journal of Bifurcation and Chaos*, vol. 25, no. 4, 2015, 1530013 (36 pages). <https://doi.org/10.1142/S021812741530013X>.
- [9] S. Hittmeyer, B. Krauskopf and H. M. Osinga. “Generalized Mandelbrot and Julia sets in a family of planar angle-doubling maps.” In M. Bohner, S. Siegmund, R. Š. Hilscher and P. Stehlík (Eds.) *Difference Equations and Discrete Dynamical Systems with Applications, ICDEA 2018*, Springer Proceedings in Mathematics and Statistics, Springer-Verlag (in press).
- [10] B. Krauskopf and H. Kriete. “Approximating transcendental Julia sets.” In H. Kriete (Ed.) *Progress in Holomorphic Dynamics*, Pitman Research Notes in Mathematics Series 387, Longman, 1998, pp. 116–138.
- [11] J. C. Mayer. “An explosion point for the set of endpoints of the Julia set of $\lambda \exp(z)$ ” *Ergodic Theory & Dynamical Systems*, vol. 10, no. 1, 1990, pp. 177–183. <https://doi.org/10.1017/S0143385700005460>.
- [12] H. M. Osinga, B. Krauskopf and S. Hittmeyer. “Chaos and wild chaos in Lorenz-type systems.” In Z. Al Sharawi, J. Cushing, and S. Elaydi (Eds.) *Theory and Applications of Difference Equations and Discrete Dynamical Systems*, Springer-Verlag, 2014, pp. 75–98.