VERY REGULAR ZERO SETS FOR THE BERGMAN SPACES

N.F. DUDLEY WARD AND P.C. FENTON

ABSTRACT. We obtain a condition which is both necessary and sufficient such that a sequence of regularly spaced points in the unit disc is a zero set for a Bergman space L_a^p.

1. Introduction

Let $\mathbb{D} = \{z : |z| < 1\}$ denote the unit disc in the complex plane and $L_a^p(\mathbb{D})$, $p \geq 1$, be the Bergman space of functions $f(z)$ analytic in $\mathbb{D}$ such that

$$\|f\|_p^p = \iint_{\mathbb{D}} |f(z)|^p dA(z) < \infty,$$

where $dA(z)$ is normalised Lebesgue area measure on $\mathbb{D}$. A sequence of points $\{z_n\}$ in $\mathbb{D}$ is called a *zero set* for L_a^p if there exists $f \not\equiv 0$ which vanishes precisely on the z_n. It is an open problem to give a geometric classification of the zeros sets for the Bergman spaces [2].

The case for the Hardy spaces H^p is simpler [1]. A sequence $\{z_n\} \subset \mathbb{D}$ is a *Blaschke sequence* provided $\sum_{n=1}^{\infty}(1 - |z_n|) < \infty$, and the zero sets for H^p are precisely the Blaschke sequences. Functions which vanish on the z_n are given by the Blaschke products:

$$B(z) = \prod_{n=1}^{\infty} \frac{z - a_n}{1 - \overline{a}_n z} \frac{-|a_n|}{a_n}.$$

All H^p spaces have the same zero sets and a union of two zero sets is a zero set. Horowitz [3] showed that neither statement holds for the Bergman spaces.

For Bergman spaces we consider zero sets of the following form. Given an integer q, $q \geq 2$, and a positive number K, let $r_j = 1 - Kq^{-j}$, for $j \geq j_0 = \log K/\log q$, and on each circle $|z| = r_j$ take $\eta_j = q^j$ equally spaced points $z_{j,1}, z_{j,2}, \ldots, z_{j,\eta_j}$:

$$z_{j,k} = r_j e^{i\theta_{j,k}}, \quad k = 1, \ldots, \eta_j, \tag{1}$$

where $\theta_{j,k} = 2\pi ki/\eta_j$. We ask for conditions on q and K that make $A = \{z_{j,k}\}$ a zero set for some Bergman space $L_a^p(\mathbb{D})$.

A case to bear in mind is the following. With $q = 2$ and $K = 1$,

Date: September 27, 2002.

Key words and phrases. Bergman spaces, zero sets.

$$z_{j,k} = \left(1 - 2^{-j}\right) e^{2\pi k i/\eta_j}, \quad k = 1, \ldots, \eta_j. \tag{2}$$

Such a set is separated in the pseudo-hyperbolic metric but is thick in the sense that a union of pseudo-hyperbolic balls with centres $z_{j,k}$ covers the disc [7]. It turns out that this set is not a zero set for L^1_a although it is a zero set for L^p_a for p sufficiently small. By thinning the sequence we obtain zero sets for larger values of the parameter p.

We prove the following result:

Theorem 1. *Given an integer q, with $q \geq 2$, and a positive number K, let $r_j = 1 - Kq^{-j}$ and $\eta_j = q^j$, $j \geq j_0 = \log K/\log q$. Suppose that $A = \{z_{j,k}\}$ is the set of points defined by (1). Then A is a zero set for L^p_a if, and only if,*

$$p < \frac{\log q}{K}.$$

The sequences defined by (1) do not satisfy the Blaschke condition. However

$$\sum_{j,k} (1 - |z_{j,k}|)^2 < \infty,$$

and so the Weierstrass product

$$\Pi(z) = \prod_{j,k} \frac{z - z_{j,k}}{1 - \overline{z}_{j,k} z} \frac{-|z_{j,k}|}{z_{j,k}} e^{\left(1 + \frac{z - z_{j,k}}{1 - \overline{z}_{j,k} z} \frac{|z_{j,k}|}{z_{j,k}}\right)},$$

converges uniformly on compact subsets of $\mathbb{D}$ and vanishes precisely on the $z_{j,k}$ [8]. Part of the proof of Theorem 1 depends on estimates of the Bergman norm of a function related to $\Pi(z)$.

2. Proof of Theorem 1

2.1. **Necessity.** Suppose that $p \geq \frac{\log q}{K}$. We will show that, given $f(z)$, analytic in $\mathbb{D}$ and having A as its zero set, $\|f\|_p^p = +\infty$.

Let $n(t)$ be the counting function of zeros of $f(z)$ and

$$N(r) = \int_0^r \frac{n(t)}{t} dt.$$

According to Jensen's formula,

$$\log |f(0)| + N(r) = \frac{1}{2\pi} \int_0^{2\pi} \log |f(re^{i\theta})| d\theta, \tag{3}$$

so that, multiplying both sides by $p > 0$ and exponentiating,

$$|f(0)|^p e^{pN(r)} = e^{\frac{1}{2\pi} \int_0^{2\pi} \log |f(re^{i\theta})|^p d\theta}.$$

Therefore, by Jensen's inequality,

$$|f(0)|^p e^{pN(r)} \leq \frac{1}{2\pi}\int_0^{2\pi} |f(re^{i\theta})|^p d\theta. \tag{4}$$

To show that $\|f\|_p^p = +\infty$ then, it is enough to show that $\int^1 re^{pN(r)}dr = +\infty$, or, what amounts to the same thing, $\int^1 e^{pN(r)}dr = +\infty$.

For $r_n \leq r < r_{n+1}$ we have

$$N(r) = \log r \sum_{j=j_0}^{n} \eta_j - \sum_{j=j_0}^{n} \eta_j \log r_j,$$

so that, since

$$r_j{}^{\eta^j} = (1 - Kq^{-j})^{q^j} < e^{-K}, \tag{5}$$

for all j, and $\log r \sum_{j=j_0}^{n} \eta_j = O(q^{n+1} \log r_{n+1} = O(1))$, we have

$$N(r) > Kn + O(1),$$

as $r \to 1$. It follows that

$$\int_{r_n}^{r_{n+1}} e^{pN(r)}dr > K(1-q^{-1})q^{-n}e^{pKn+O(1)} \geq K(1-q^{-1})e^{O(1)},$$

and therefore $\int^1 e^{pN(r)}dr = +\infty$.

2.2. **Sufficiency.** We will show if $p < (\log q)/K$, then, with j_0 as in the statement of the theorem,

$$f(z) = \prod_{j=j_0}^{\infty} \pi_j(z)\sigma_j(z)$$

belongs to L_a^p, where

$$\pi_j(z) = \prod_{k=1}^{\eta_j} \frac{z - z_{j,k}}{1 - \overline{z}_{j,k}z} \frac{-|z_{j,k}|}{z_{j,k}} e^{\left(1 + \frac{z - z_{j,k}}{1 - \overline{z}_{j,k}z} \frac{|z_{j,k}|}{z_{j,k}}\right)}$$

and

$$\sigma_j(z) = \exp\left(-\frac{2Kr_j^{\eta_j} z^{\eta_j}}{1 - r_j^{\eta_j} z^{\eta_j}}\right).$$

Evidently $f(z)$ vanishes precisely at the points of A, so that A is a zero set for L_a^p.

Consider first $\pi_j(z)$. The general term in $\pi_j(z)$ has the form $(1-u)e^u$, where

$$u = 1 + \left(\frac{z - z_{j,k}}{1 - \overline{z}_{j,k}z}\right)\frac{|z_{j,k}|}{z_{j,k}} = (1 + z|z_{j,k}|/z_{j,k})\frac{1 - |z_{j,k}|}{(1 - \overline{z}_{j,k}z)},$$

and since $|(1-u)e^u| \leq e^{|u|^2/2}$, $|u| \leq 2$, we have

$$|\pi_j(z)| \leq e^{2\sum_{k=1}^{\eta_j}\left(\frac{1-|z_{j,k}|}{|1-\overline{z}_{j,k}z|}\right)^2} = e^{2\sum_{k=1}^{\eta_j}\left(\frac{1-r_j}{|1-\overline{z}_{j,k}z|}\right)^2}.$$

Now, with $z = re^{i\theta}$,

$$\begin{aligned}\left(\frac{1-r_j}{|1-\overline{z}_{j,k}z|}\right)^2 &= \frac{(1-r_j)^2}{1-r_j^2r^2}\frac{1-r_j^2r^2}{|1-\overline{z}_{j,k}z|^2}\\ &= \frac{(1-r_j)^2}{1-r_j^2r^2}\sum_{m=-\infty}^{\infty} r_j^{|m|}r^{|m|}e^{im(\theta_{j,k}-\theta)}.\end{aligned}$$

Also $\theta_{j,k} = 2\pi k q^{-j}$, $k = 1, 2, \ldots, \eta_j = q^j$ are the η_jth roots of unity and so for m not an integer multiple of η_j, $\sum_{k=1}^{\eta_j} e^{im\theta_{j,k}} = 0$. For $m = l\eta_j$

$$\sum_{k=1}^{\eta_j} e^{im\theta_{j,k}} = \eta_j = q^j.$$

Therefore

$$\begin{aligned}\sum_{k=1}^{\eta_j}\left(\frac{1-r_j}{|1-\overline{z}_{j,k}z|}\right)^2 &= \frac{K(1-r_j)}{1-r_j^2r^2}\sum_{l=-\infty}^{\infty} r_j^{|l|\eta_j}r^{|l|\eta_j}e^{-il\eta j\theta}\\ &= \frac{K(1-r_j)}{1-r_j^2r^2}\frac{1-r_j^{2\eta_j}r^{2\eta_j}}{|1-r_j^{\eta_j}z^{\eta_j}|^2}\\ &= \frac{K(1-r_j)}{1-r_j^2r^2}\Re\left(\frac{1+r_j^{\eta_j}z^{\eta_j}}{1-r_j^{\eta_j}z^{\eta_j}}\right),\end{aligned}$$

and so

$$|\pi_j(z)| \le \exp\left(\frac{2K(1-r_j)}{1-r_j^2r^2}\Re\left(\frac{1+r_j^{\eta_j}z^{\eta_j}}{1-r_j^{\eta_j}z^{\eta_j}}\right)\right).$$

Supposing that $r_n \le r < r_{n+1}$, we consider two cases: j satisfying $1 \le j \le n$, and j satisfying $j \ge n+1$.

(i) $j \ge n+1$. Since

$$\frac{1-r_j}{1-r_j^2r^2} \le \frac{1-r_j}{1-r^2} \le \frac{1-r_j}{1-r} \le \frac{1-r_j}{1-r_{n+1}} = q^{n+1-j},$$

we have, taking account of (5),

$$\begin{aligned}|\pi_j(z)| &\le \exp\left(2q^{n+1-j}\left(\frac{1+r_j^{\eta_j}r^{\eta_j}}{1-r_j^{\eta_j}r^{\eta_j}}\right)\right)\\ &\le \exp\left(2q^{n+1-j}\left(\frac{1+r_j^{\eta_j}}{1-r_j^{\eta_j}}\right)\right)\\ &\le \exp\left(2q^{n+1-j}\frac{e^K+1}{e^K-1}\right),\end{aligned}$$

and therefore, for all large n,

$$\prod_{j=n+1}^{\infty} |\pi_j(z)| \leq \exp\left(\frac{2(e^K+1)}{(1-q^{-1})(e^K-1)}\right).$$

(ii) $j_0 \leq j \leq n$. Since

$$\frac{1-r_j}{1-r_j^2 r^2} \leq \frac{1-r_j}{1-r_j^2} = \frac{1}{1+r_j} = \frac{1}{2} + \frac{Kq^{-j}}{2-Kq^{-j}} < \frac{1}{2} + Kq^{-j}/2,$$

and

$$\Re\left(\frac{1+r_j^{\eta_j} z^{\eta_j}}{1-r_j^{\eta_j} z^{\eta_j}}\right) = 1 + \Re\left(\frac{2r_j^{\eta_j} z^{\eta_j}}{1-r_j^{\eta_j} z^{\eta_j}}\right),$$

and also, from (5),

$$\Re\left(\frac{2r_j^{\eta_j} z^{\eta_j}}{1-r_j^{\eta_j} z^{\eta_j}}\right) \leq \frac{2}{1-e^{-K}},$$

we have

$$|\pi_j(z)| \leq \exp\left(K + \Re\left(\frac{2Kr_j^{\eta_j} z^{\eta_j}}{1-r_j^{\eta_j} z^{\eta_j}}\right) + \frac{3K^2 q^{-j}}{1-e^{-K}}\right).$$

Recalling the definition of $\sigma_j(z)$, then,

$$\prod_{j=1}^{n} |\pi_j(z)\sigma_j(z)| \leq \exp\left(Kn + \frac{3K^2}{(1-q^{-1})(1-e^{-K})}\right).$$

Finally

$$\begin{aligned}
\prod_{j=n+1}^{\infty} |\sigma_j(z)| &\leq \exp\left(\frac{2Ke^{-K}}{1-e^{-K}} \sum_{j=n+1}^{\infty} r^{\eta_j}\right) \\
&\leq \exp\left(\frac{2Ke^{-K}}{1-e^{-K}} \sum_{j=n+1}^{\infty} (1-Kq^{-n-1})^{q^{n+1}Kq^{j-n-1}}\right) \\
&\leq \exp\left(\frac{2Ke^{-K}}{1-e^{-K}} \sum_{j=0}^{\infty} e^{-K^2 q^j}\right),
\end{aligned}$$

and, since $q^j > (q-1)j$ for all $j \geq 0$, this last series is convergent,and thus $\prod_{j \geq n+1} |\sigma_j(z)|$ is bounded independently of z.

Combining our results we obtain, for $r_n \leq r < r_{n+1}$, $|f(z)| \leq Me^{Kn}$, where $M = M(q,K)$ is a constant. Hence

$$\begin{aligned}
\int_{r_n \leq r < r_{n+1}} |f(z)|^p dA(z) &\leq 2\pi K M^p e^{Kpn}(q^{-n} - q^{-n-1}) \\
&= 2\pi K M^p (1-q^{-1}) e^{(Kp - \log q)n}.
\end{aligned}$$

Since $Kp < \log q$,

$$\int_{r_1 \le r < r_{n+1}} |f(z)|^p dA(z) = \sum_{n=1}^{\infty} \int_{r_n \le r < r_{n+1}} |f(z)|^p dA(z) < \infty,$$

and $f \in L^p_a$.

Remarks . (i) Let $r_j = 1 - K2^{-j}$. For $K = 1$, $A = \{z_{j,k}\}$ is a zero set for L^p_a with $p < 0.693\ldots$. For $K = 1/2$, A is a zero set for $p < 1.386\ldots$ and for $K = 1/4$, for $p < 2.772\ldots$.

(ii) We obtain easily zeros sets $A = \{z_{j,k}\}$ and $B = \{w_{j,k}\}$ whose union is not a zero set for the same L^p_a. For example take

$$\begin{aligned} z_{j,k} &= \left(1 - 1/2^j\right) e^{2\pi k i/2^j}, \\ w_{j,k} &= \left(1 - 1/2^j\right) e^{2\pi(k+1/2)i/2^j}. \end{aligned}$$

(iii) Plainly in order to obtain a geometric description of more general zero sets one has to consider also the angular distribution of the zeros. In two seminal papers [4], [5] Korenblum uses the Beurling-Carleson characteristic to give a geometric classification of certain growth spaces related to the Bergman spaces. In particular he characterised the zero sets for the union of all Bergman spaces.

In conclusion we express our warm gratitude to Professors Milne Anderson and Walter Hayman FRS for discussions which lead to this work. We are also grateful to Professor Korenblum for drawing our attention to a paper of Professor Leucking [6] who also proves Theorem 1.

References

[1] J. B. Garnett. *Bounded Analytic Functions.* Academic Press, 1981.
[2] H. Hedenmalm, B. Korenblum, and K. Zhu. *Theory of Bergman Spaces.* Springer, 2000.
[3] C. Horowitz. Zeros of functions in the Bergman spaces. *Duke Math. J.*, 41:693–710, 1974.
[4] B. Korenblum. An extension of the Nevanlinna theory. *Acta Mathematica*, 135:187–219, 1975.
[5] B. Korenblum. A Beurling-type theorem. *Acta Mathematica*, 138:265–293, 1976.
[6] D. H. Luecking. Zero sequences for Bergman spaces. *Complex Variables*, 30:345–362, 1996.
[7] E. M. Stein. *Singular integrals and differentiability properties of functions.* Princeton, 1970.
[8] E. C. Titchmarsh. *Theory of Functions.* Oxford University Press, 1954.

Department of Mathematics, University College, Gower Street, London, WC1E 6BT, UNITED KINGDOM.

Current address: Department of Mathematics, Auckland University, Private Bag 92019, Auckland, NEW ZEALAND.

E-mail address: nfdw@math.ucl.ac.uk

Department of Mathematics, Otago University, PO Box 56, Dunedin, NEW ZEALAND.

E-mail address: pfenton@maths.otago.ac.nz