

# Fitting Parameters for a Solvable Model of a Quantum Network

M. Harmer

email: harmer@math.auckland.ac.nz

Department of Mathematics

University of Auckland

New Zealand

April 26, 2004

## Abstract

A solvable model corresponding to a given quantum network is described in [21] without an explicit description of how to fit the parameters of the solvable model. Here we give a procedure to fit these parameters so that the solvable model reproduces the important features, viz. the scattering matrix for the physically relevant energies, of the quantum network, subject to the non-vanishing of a determinant.

## 1 Introduction

The current interest in quantum networks is motivated by the design of electronic devices at quantum scales with the objective of manufacturing networks with prescribed transport properties. The literature in this area is extensive with many studies into the mathematical aspects of such problems [1, 2, 5, 6, 7, 8, 11, 14, 15, 12, 13, 17, 18, 19, 21, 23, 25, 24, 22, 26] as well as explicit attempts at the actual design of networks with given scattering properties [3, 4, 9, 10, 16, 20, 27]. We do not make any claim for completeness of our bibliography.

In [21] the authors describe how to associate with a given quantum network a solvable model with the same scattering properties at physically relevant energies. However, it is not shown that the desired solvable model exists, rather some conditions are found which this solvable model has to satisfy. Here we show (subject to the non-vanishing of a determinant) that the desired solvable model can always be constructed and give the explicit construction. We start with a brief description of the quantum network and associated solvable model proposed in [21]. Our notation is the same as, or a simplified version

of, the notation in [21].

## 2 The Quantum Network

The quantum network consists of a connected set of compact domains  $\{\Omega_i\}$  and finite and semi-infinite channels  $\{\omega^j\}$  lying in the plane  $\Omega = \bigcup_i \Omega_i \bigcup_j \omega^j \subset \mathbb{R}^2$ . The channel  $\omega^j$  has straight edges and constant width  $\delta_j$  and is connected to one compact domain (if it is semi-infinite) or two compact domains (if it is finite). The boundary arc  $\gamma_i^j$  between  $\Omega_i$  and  $\omega^j$  is linear and at right angles to the edges of  $\omega^j$ . We denote by  $\Gamma = \bigcup_{i,j} \gamma_i^j$  the set of boundaries.

We consider the spectral problem on the network

$$\begin{aligned} \mathcal{L}\psi &\equiv -\Delta\psi + V\psi = \lambda\psi \\ \psi|_{\partial\Omega} &= 0 \end{aligned}$$

where the potential  $V : \Omega \mapsto \mathbb{R}$  is real and assumed constant on the channels  $V|_{\omega^j} \equiv V_j = \text{const}$ .

By the Glazman splitting technique we see that the continuous spectrum, regardless of the form of the compact part of the network, consists of an infinite union of semi-infinite intervals, one for each mode transverse to the channel, and of multiplicity equal to the number of semi-infinite channels. It is assumed that the Fermi energy  $\Lambda$  is in the continuous spectrum. The authors of [21] define the ‘intermediate operator’  $\mathcal{L}_\Lambda$  in terms of special boundary conditions at  $\Gamma$  which depend on the Fermi energy.

Then the Dirichlet to Neumann map for this intermediate operator  $\mathcal{DN}^\Lambda(\lambda)$  is calculated and it is shown that the scattering matrix is given in terms of this operator by ([21], equation (24))

$$\mathbf{S}(\lambda) = - \left( \mathcal{DN}^\Lambda - \frac{i}{2\mu^\parallel} K_+ \right)^{-1} \left( \mathcal{DN}^\Lambda + \frac{i}{2\mu^\parallel} K_+ \right). \quad (1)$$

Here  $\mu^\parallel$  is a constant and  $K_+$  is a simple function of the spectral parameter. For the purposes of approximation by a solvable model the authors use the so called ‘essential part’  $\mathcal{DN}_T^\Lambda$  of the Dirichlet to Neumann map obtained by only considering eigenvalues in a band around the Fermi energy of width proportional to the temperature. This operator has kernel ([21], equation (48), although there is some discrepancy with equation (35))

$$\mathcal{DN}_T^\Lambda = \sum_{l=1}^{N_T} \frac{1}{\lambda_l - \lambda} \left| P_+ \frac{\partial\varphi_l}{\partial n} \right\rangle \left\langle P_+ \frac{\partial\varphi_l}{\partial n} \right|.$$

Here the  $\varphi_l$  and  $\lambda_l$  are the eigenfunctions and eigenvalues on the compact part of the network,  $\partial \cdot / \partial n$  is the normal derivative and  $P_+$  is a projection onto a finite dimensional subspace on  $\Gamma$ , related to the finite multiplicity of

the continuous spectrum at the Fermi energy. Consequently, this ‘essential part’ of the Dirichlet to Neumann operator is a finite dimensional operator, ie. a matrix.

Consider an approximation to the scattering matrix obtained by using this simpler operator  $\mathcal{DN}_T^\Lambda$  in (1). One of the aims of [21] is to construct a solvable model which has the same scattering matrix as this approximate scattering matrix. Some conditions are found for this construction to proceed but it is not shown that it is always possible to construct the desired solvable model. Here we will show (modulo the non-vanishing of a determinant) that it is always possible to construct a solvable model with the desired properties.

### 3 The Solvable Model and Fitting Parameters

In [21] the authors approximate the quantum network by a (one-dimensional) graph consisting of  $n$  semi-infinite rays and a single vertex with a finite dimensional operator  $A$  at the vertex. The set of finite dimensional vectors and eigenvalues

$$\left\{ P_+ \frac{\partial \varphi_l}{\partial n}, \lambda_l; l = 1, \dots, N_T \right\} \quad (2)$$

are regarded as given data from the quantum network. Following [21] the vectors  $P_+ \partial \varphi_l / \partial n$  are defined to be elements of the  $n$ -dimensional<sup>1</sup> vector space  $E_+$ .

We denote by  $\ell = \bigoplus^n -\frac{d^2}{dx^2}$  the ‘Schrödinger’ operator on the rays of the graph acting on the Hilbert space  $H = \bigoplus^n L^2[0, \infty)$ . We denote by  $E_A = \mathbb{C}^N$  the ‘abstract’ finite dimensional Hilbert space at the vertex on which the self-adjoint matrix  $A = A^*$  acts. These are clearly both self-adjoint operators in their respective Hilbert spaces.

We restrict  $\ell$  to a *symmetric* operator  $\ell_0 = \ell|_{\mathcal{D}_0}$  acting on functions disappearing near the ‘origins’ (of the  $L^2[0, \infty)$ ).

We also restrict  $A$  to the domain  $D_0 \equiv (A - i\mathbb{I})^{-1} E_A \ominus N_i$  where  $N_i$ , the deficiency subspace with  $\dim(N_i) = d$ , is chosen in the construction of the solvable model. A technical requirement is that  $N_{-i} \cap N_i = \emptyset$  where

$$N_{-i} = \frac{A + i}{A - i} N_i,$$

ie. the deficiency subspaces do not intersect. For this to hold we need at least  $2d \leq N$ .

Our aim is to find a self-adjoint extension of the symmetric operator

$$\mathbf{A}_0 = \ell_0 \oplus A_0 \quad (3)$$

---

<sup>1</sup>From [21] we see that the dimension of these vectors is *at least* the number of semi-infinite channels of the network.

which has the same scattering matrix as the network in the desired energy range.

This procedure of restricting then extending the operator to a self-adjoint operator with desired properties is described in some detail in [26, 21]. In summary the scattering ansatz for the solvable model ([21], after equation (43)) takes the form

$$\Psi = \begin{pmatrix} e^{-iK+\mathbf{x}}\nu + e^{iK+\mathbf{x}}\mathbf{S}\nu \\ \frac{A+i}{A-\lambda}\xi \end{pmatrix}$$

where  $\xi$  are the components of  $\Psi$  in the deficiency subspace. We denote by  $\beta$  the self-adjoint boundary conditions for the extension  $\mathbf{A}_\beta$  of (3) (to be precise, in this report  $\beta$  is used to denote the off diagonal part of the boundary condition matrix). These boundary conditions relate  $\Psi$  and its derivative at the origin allowing the authors of [21] to solve for the scattering matrix in terms of  $\beta$ ,  $A$  and the deficiency subspaces. This gives theorem 5.1 of [21] which describes the solvable model in terms of the set (2) of given data from the quantum network. Our formulation of this theorem follows the formulation in [21] closely although we have omitted the last two sentences as they are not strictly relevant to our discussion.

**Theorem 5.1** *The constructed operator  $\mathbf{A}_\beta$  is a solvable model of the quantum network on the essential interval of energy iff: the dimension of the space  $E_A$  coincides with the number of eigenvalues on the essential interval of energy  $N = N_T$ ; the eigenvalues  $\lambda_l$  of  $A$  coincide with the eigenvalues of the intermediate operator on the essential interval of energy  $\lambda_l = \lambda_l$  so that*

$$A = \sum_{l=1}^{N_T} \lambda_l |e_l\rangle \langle e_l| ;$$

*and there exists a deficiency subspace  $N_i$  of  $A_0$ , the associated orthogonal projection  $P_i$  onto  $N_i$ , and an operator  $\beta : N_i \mapsto E_+$  such that for the orthonormal basis of eigenvectors  $e_l$  of  $A$*

$$P_+ \frac{\partial \varphi_l}{\partial n} = \beta P_i e_l \tag{4}$$

*for  $l = 1, \dots, N_T$ .*

**Note:** the matrix  $\beta$ , although associated with the self-adjoint boundary conditions, is not necessarily self-adjoint because it is the off diagonal block of the boundary condition matrix.

**Note:** as noted above the dimension of  $N_i$  has to be less than or equal half the dimension of  $E_A$ ,  $2d \leq N_T$ , otherwise there is no possibility that the intersection of the deficiency subspaces is null. The condition of null intersection is expressed below as the non-vanishing of a determinant; however, if

this determinant vanishes our procedure fails<sup>2</sup>.

Really, this theorem defines the solvable model up to the condition (4). We show that it is always possible to choose  $P_i$  (and thence  $N_i$ ) and  $\beta$  to satisfy this condition. We start by defining the  $n$ -dimensional vector

$$\Phi_l = P_+ \frac{\partial \varphi_l}{\partial n}.$$

Considering the  $\Phi_l$  as columns of a  $n \times N_T$  matrix  $\Phi$  we can write the condition (4) in matrix form  $\Phi = \beta P_i$  where we have implicitly assumed that  $\{e_l\}$  is the standard basis (so  $A$  is a diagonal matrix in this basis). We diagonalise the self-adjoint matrix

$$\Phi \Phi^* = U^* D U \tag{5}$$

and define the  $n \times N_T$  matrix

$$\Upsilon = U \Phi. \tag{6}$$

Consequently condition (4) becomes

$$\Upsilon = U \beta P_i \tag{7}$$

and we claim that this has solution

$$\beta = U^* D^{\frac{1}{2}} \tag{8}$$

$$P_i = D^{-\frac{1}{2}} \Upsilon \tag{9}$$

where we take the real non-negative square root of  $D$ —it follows easily from (5) that the components of  $D$  are non-negative. When  $D$  has zeroes on the diagonal we make sense of (9) by defining  $D^{-\frac{1}{2}}$  to have zeroes at the corresponding positions in the diagonal.

Clearly (8,9) formally satisfies (4) or (7) and we will show that this holds even when  $D$  has zeroes. As we pointed out above  $\beta$  can take any form so the choice of  $\beta$  needs no further discussion. We just need to show that  $P_i$  is an orthogonal projection for suitable bases.

Assuming for now that  $D$  has no zeroes we consider following set

$$\eta_k = \sum_{l=1}^{N_T} \left( D^{-\frac{1}{2}} \Upsilon \right)_{k,l} e_l \tag{10}$$

of vectors in  $E_A$ . Since  $\{e_l\}$  is orthonormal, and since (5) and (6) give us

$$\Upsilon \Upsilon^* = D, \tag{11}$$

we see that  $\{\eta_k\}_{k=1}^n$  is also an orthonormal set in  $E_A$ . Now we claim that this is the basis of  $N_i$ . In consequence

$$P_i = \sum_{k=1}^n |\eta_k\rangle \langle \eta_k|,$$

---

<sup>2</sup>Although it appears that the condition of null-intersection may be relaxed.

with domain in the basis  $\{e_l\}$  and range in the basis  $\{\eta_k\}$ , just takes the desired form (9). We note that in this case ( $D$  having no zeroes) the deficiency index is  $d = n$ .

In the case where  $D$  has zeroes on the diagonal we easily see from (11) that the corresponding rows of  $\Upsilon$  are *identically zero*. Consequently, using the given definition of  $D^{-\frac{1}{2}}$  we see that  $D^{\frac{1}{2}}D^{-\frac{1}{2}}$  has ones on the diagonal except at those positions corresponding to the zero rows of  $\Upsilon$  so that

$$D^{\frac{1}{2}}D^{-\frac{1}{2}}\Upsilon = \Upsilon$$

and (8,9) continues to satisfy (4), (7). Furthermore the definition of the basis (10) continues to make sense as does the demonstration that  $P_i$  is an orthogonal projection. We note that in this case the deficiency index is  $d = \text{rank}(D) < n$ .

The condition for the non-intersection  $N_{-i} \cap N_i = \emptyset$  can be written in terms of the non-vanishing of a determinant involving  $D^{-\frac{1}{2}}\Upsilon$  and the eigenvalues  $\lambda_l$ . Really, working in the basis  $\{e_l\}$  of  $E_A$  we see that  $N_i$  has orthonormal basis given by the rows of  $D^{-\frac{1}{2}}\Upsilon$ . Since the Cayley transform  $(A + i)(A - i)^{-1}$  is unitary,  $N_{-i}$  has orthonormal basis given by the rows of  $D^{-\frac{1}{2}}\Upsilon \text{diag}((\lambda_l + i)/(\lambda_l - i))$ . Consequently, the deficiency subspaces have null intersection iff

$$\det \begin{pmatrix} D^{-\frac{1}{2}}\Upsilon \\ D^{-\frac{1}{2}}\Upsilon \text{diag} \left( \frac{\lambda_l + i}{\lambda_l - i} \right) \end{pmatrix} \neq 0 \quad (12)$$

where of course we omit any zero rows in the matrix  $D^{-\frac{1}{2}}\Upsilon$ .

In summary, given the data (2) we find  $D$ ,  $U$  and  $\Upsilon$  from equations (5) and (6). The boundary condition for the desired solvable model is then given by equation (8) and the deficiency subspace is given by (9,10) where the deficiency index is  $d = \text{rank}(D) \leq n$ . We note that there are no free parameters to ensure that the condition for the null intersection of the deficiency subspaces (12) is satisfied.

## References

- [1] S. Albeverio, F. Gesztesy, R. Hoegh-Krohn, and H. Holden. *Solvable models in quantum mechanics*. Springer-Verlag, New York, 1988.
- [2] S. Albeverio and P. Kurasov. *Singular Perturbations of Differential Operators*. London Mathematical Society Lecture Note Series 271. Cambridge University Press, Cambridge, 2000.
- [3] I. Antoniou, B. S. Pavlov, and A. M. Yafyasov. Quantum electronic devices based on metal-dielectric transition in low-dimensional quantum

- structures. In D. S. Bridges, C. Calude, J. Gibbons, S. Reeves, and I. Witten, editors, *Combinatorics, Complexity, Logic (Proceedings of DMTCS '96)*, Singapore, 1996. Springer.
- [4] V. Bogevolnov, A. Mikhailova, B. S. Pavlov, and A. Yafyasov. About scattering on the ring. Department of Maths Report series 413 ISSN 1173-0889, University of Auckland, New Zealand, 1999. To be published in ‘Operator Theory: Advances and Applications’, (IWOTA 98, dedicated to I. Gohberg).
- [5] W. Bulla and T. Trenkler. The free dirac operator on compact and noncompact graphs. *J Math. Phys.*, 31:1157–1163, 1990.
- [6] R. Carlson. The second derivative operator for a weighted graph. preprint.
- [7] P. Exner. Contact interactions on graph superlattices. *J. Phys. A: Math. Gen.*, 29:87–102, 1996.
- [8] P. Exner and P. Šeba. Free quantum motion on a branching graph. *Rep. Math. Phys.*, 28:7–26, 1989.
- [9] P. Exner, P. Šeba, and P. Stovicek. Quantum interference on graphs controlled by an external electrical field. *J. Phys. A: Math. Gen.*, 21:4009–4019, 1988.
- [10] B. Gejler, B. S. Pavlov, and I. Popov. Possible construction of a quantum triple logic device. New technologies for narrow-gap semiconductors, progress report (july 1, 1998–june 30 1999), ESPRIT NTCONGS, 1999. ESPRIT project No. 28890.
- [11] N. I. Gerasimenko. The inverse scattering problem on a noncompact graph. *Teoret. Mat. Fiz.*, 75:187–200, 1988.
- [12] M. Harmer. Hermitian symplectic geometry and extension theory. *Journal of Physics A: Mathematical and General*, 33:9193–9203, 2000.
- [13] M. Harmer. Hermitian symplectic geometry and the factorisation of the scattering matrix on graphs. *Journal of Physics A: Mathematical and General*, 33:9015–9032, 2000.
- [14] M. Harmer. *The Matrix Schrödinger Operator and Schrödinger Operator on Graphs*. PhD thesis, University of Auckland, 2000.
- [15] M. Harmer. Inverse scattering for the matrix Schrödinger operator and Schrödinger operator on graphs with general self-adjoint boundary conditions. *ANZIAM Journal*, 43:1–8, 2002.

- [16] M. Harmer, B. S. Pavlov, and A. Mikhailova. Manipulating the electron current through a splitting. *Proceedings of the Centre for Mathematics and its Applications*, 39:118–131, 2001.
- [17] V. Kostykin and R. Schrader. Kirchhoff’s rule for quantum wires. *J. Phys A: Math. Gen.*, 32:595–630, 1999.
- [18] Yu. B. Melnikov. Scattering on graphs as a quantum few-body problem. preprint IPRT #09-93, 1993.
- [19] Yu. B. Melnikov and B. S. Pavlov. Two-body scattering on a graph and application to simple nanoelectronic devices. *J. Math. Phys.*, 36:2813–2825, 1995.
- [20] A. Mikhailova, B. S. Pavlov, I. Popov, T. Rudakova, and A. M. Yafyasov. Scattering on a compact domain with few semi-infinite wires attached: resonance case. Department of Maths Report series 420, University of Auckland, New Zealand, 1999. submitted to *Mathematische Nachrichten*.
- [21] A. B. Mikhailova, B. S. Pavlov, and L. V. Prokhorov. Modeling of quantum networks. unpublished.
- [22] S. P. Novikov. Discrete schrödinger operators and topology. preprint, dedicated to the 70-th birthday of Mikio Sato.
- [23] S. P. Novikov. Schrödinger operators on graphs and topology. *Uspekhi Math. Nauk-Russian. Math. Surveys*, 52(6):127–168, 1997.
- [24] S. P. Novikov. Schrödinger operators on graphs and symplectic geometry. In E. Bierstone, B. Khesin, A. Khovanskii, and J. Marsden, editors, *The Arnol’dfest*, 1998. Proceedings of the Fields Institute Conference in Honour of the 60-th Birthday of Vladimir I. Arnol’d, to appear in the Fields Institute Communications Series.
- [25] S. P. Novikov and I. A. Dynnikov. Discrete spectral symmetries of low-dimensional differential operators and difference operators on regular lattices and two-dimensional manifolds. *Uspekhi Math. Nauk-Russian. Math. Surveys*, 52(5):1057–1116, 1997.
- [26] B. S. Pavlov. The theory of extensions and explicitly solvable models. *Uspekhi Math. Nauk-Russian. Math. Surveys*, 42(6):127–168, 1987.
- [27] A. M. Yafyasov, V. B. Bogevolnov, and T. V. Rudakova. Quantum interferential electronic transistor (qiet). theoretical analysis of electronic properties for low-dimensional systems. (Progress Report 1995), Preprint IPRT N 99-95.