

On adding a list of numbers (and other one-dependent determinantal processes)

Alexei Borodin* Persi Diaconis† Jason Fulman‡

Version of April 6, 2009

Abstract

Adding a column of numbers produces “carries” along the way. We show that random digits produce a pattern of carries with a neat probabilistic description: the carries form a one-dependent determinantal point process. This makes it easy to answer natural questions: How many carries are typical? Where are they located? We show that many further examples, from combinatorics, algebra and group theory, have essentially the same neat formulae, and that any one-dependent point process on the integers is determinantal. The examples give a gentle introduction to the emerging fields of one-dependent and determinantal point processes.

1 Introduction

Consider the task of adding a single column of digits:

$$\begin{array}{r} 7 \cdot 7 \\ 9 \cdot 6 \\ 4 \cdot 0 \\ 8 \cdot 8 \\ 3 \cdot 1 \\ 6 \cdot 7 \\ 1 \cdot 8 \\ 6 \cdot 4 \\ 8 \cdot 2 \\ \hline 52 \end{array}$$

We have put a dot to the right of a digit whenever the succeeding addition leads to a carry. The remainder (mod 10) is written to the right of the dot. Thus $7 + 9 = 16$ results in a dot by the 7 and a remainder of 6. At the end, the total is found by adding the number of dots (here, five) and appending the final remainder (here, 2) for a total of 52. The dots are a standard bookkeeping device used (for example) in the Trachtenberg system of speed addition [16].

**Department of Mathematics, Caltech*

†*Departments of Mathematics and Statistics, Stanford*

‡*Department of Mathematics, USC*

How many carries are typical and how are they distributed? Common sense suggests that about half the places have carries and that if a carry occurs, it is less likely that the next addition gives a carry. Investigating these questions when digits are chosen uniformly at random leads to interesting mathematics.

As additional motivation, look at the remainder column in the example above. Observe that there is a carry on the left (and so a dot) if and only if there is a descent on the right (a sequence x_1, x_2, \dots has a descent at i if $x_i > x_{i+1}$). Further, if the original column of digits is independent and uniformly distributed, so is the remainder column. Thus, the results about carries give the distribution of the descent pattern in a random sequence of digits.

Section 2 derives the distribution theory of carries by elementary arguments. There are nice formulae for the chance of any pattern of carries, and the process is one-dependent so a variety of probabilistic limit theorems are available.

Section 3 introduces determinantal point processes, shows that the carries process is determinantal, and illustrates how standard tools for determinantal point processes apply to carries.

Section 4 reviews the literature on stationary one-dependent processes, shows that all of these are determinantal and further that essentially all the neat formulae for carries and descents have versions for any stationary one-dependent process. Connections with symmetric function theory are developed. A large class of examples arising from the work of Polishchuk and Positselski [58] on Koszul algebras is shown to yield many natural examples (including the original carries process).

Section 5 gives combinatorial examples: descents in permutations from both uniform and non-uniform distributions, the connectivity set of a permutation, and binomial posets.

Section 6 generalizes from adding numbers to multiplying random elements in a finite group. For central extensions (e.g., the quaternions), there is again a carries process that is stationary, one-dependent, and determinantal. Thus explicit formulae and limit theorems are available.

Section 7 contains proofs of some of our more technical results. It shows that any one-dependent (possibly non-stationary) point process on the integers is determinantal. It also constructs a family of one-dependent determinantal processes (generalizing many examples in earlier sections), and computes its correlation kernel.

2 Probability theory for carries and descents

Throughout this section we work base b and so with the alphabet $\mathcal{B} = \{0, 1, \dots, b-1\}$. Let B_1, B_2, \dots, B_n be a sequence of randomly chosen elements of \mathcal{B} (independent and identically distributed). There is a *descent at i* if $B_i > B_{i+1}$, $1 \leq i \leq n-1$. Let X_i be 1 or 0 as there is a descent at i and $D = \{i : X_i = 1\}$ be the descent set of B_1, B_2, \dots, B_n .

The following probabilistic facts are elementary. They are stated for carries but verification is easier using descents.

Fact 1. (Single carries) For any $i \in [n-1]$,

$$P(X_i = 1) = \frac{1}{2} - \frac{1}{2b} = \frac{\binom{b}{2}}{b^2}.$$

Thus, when $b = 10$, the chance of a carry is .45. When $b = 2$, the chance of a carry is .25. For any base, $\text{Var}(X_i) = \frac{1}{4} - \frac{1}{4b^2}$.

Fact 2. (Runs of carries) For any i and j with $1 \leq i < i + j \leq n$,

$$P(X_i = X_{i+1} = \cdots = X_{i+j-1} = 1) = \binom{b}{j+1} / b^{j+1}.$$

Thus, a run of b or more carries is impossible. Further,

$$\text{Cov}(X_i, X_{i+1}) = E(X_i X_{i+1}) - E(X_i)E(X_{i+1}) = -\frac{1}{12} \left(1 - \frac{1}{b^2}\right).$$

Fact 3. (Stationary one-dependence) The distribution of $\{X_i\}_{i \in [n-1]}$ is stationary: for $J \subseteq [n-1]$, $i \in [n-1]$ with $J+i \subseteq [n-1]$, the distribution of $\{X_j\}_{j \in J}$ is the same as the distribution of $\{X_j\}_{j \in J+i}$. Further, the distribution of $\{X_i\}_{i \in [n-1]}$ is one-dependent: if $J \subseteq [n-1]$ has $j_1, j_2 \in J \Rightarrow |j_1 - j_2| > 1$, then $\{X_j\}_{j \in J}$ are jointly independent binary random variables with $P(X_j = 1) = \frac{1}{2} - \frac{1}{2b}$. The literature on m -dependent random variables is extensive. In particular, a classical central limit theorem [40] shows the following:

Theorem 2.1. For $n \geq 2$, the total number of carries $T_{n-1} = X_1 + \cdots + X_{n-1}$ has mean $(n-1)(\frac{1}{2} - \frac{1}{2b})$, variance $\frac{n+1}{12}(1 - \frac{1}{b^2})$ and, normalized by its mean and variance, T_{n-1} has a standard normal limiting distribution for n large.

Remarks.

1. An $O(n^{-1/2})$ error bound in this central limit theorem can be proved using the dependency graph approach to normal approximation by Stein's method [12], [18].
2. Stationary pairwise independent processes can fail to obey the central limit theorem [7], [43].

Fact 4. (k -point correlations) For $A \subseteq [n-1]$, let

$$\rho(A) = P\{X_i = 1 \text{ for } i \in A\}.$$

For $|A| = k$, the $\rho(A)$ are called k -point correlations for the point process X_1, \dots, X_{n-1} . They are basic descriptive units for general point processes [17]. For the carries and descent process, they are simple to describe. Break $A \subseteq [n-1]$ into disjoint, non-empty blocks of consecutive integers $A = A_1 \cup A_2 \cup \cdots \cup A_k$. Thus $A = \{2, 3, 5, 6, 7, 11\} = \{2, 3\} \cup \{5, 6, 7\} \cup \{11\}$. From Facts 2 and 3 above,

$$\rho(A) = \prod_{i=1}^k \left[\binom{b}{a_i+1} / b^{a_i+1} \right] \quad \text{if } A = \bigcup_{i=1}^k A_i \text{ with } |A_i| = a_i.$$

Fact 5. (Determinant formula) Let $\epsilon_1, \epsilon_2, \dots, \epsilon_{n-1}$ be a fixed sequence in $\{0, 1\}$. Then

$$P\{X_1 = \epsilon_1, \dots, X_{n-1} = \epsilon_{n-1}\} = \frac{1}{b^n} \cdot \det \begin{pmatrix} s_{j+1} - s_i + b - 1 \\ b - 1 \end{pmatrix}.$$

Here, if there are exactly k 1's in the ϵ -sequence at positions $s_1 < s_2 < \cdots < s_k$, the determinant is of a $(k+1) \times (k+1)$ matrix with (i, j) entry $\binom{s_{j+1} - s_i + b - 1}{b - 1}$ for $0 \leq i, j \leq k$ with $s_0 = 0$, $s_{k+1} = n$.

Example 1. If $n = 8$, $\epsilon_1 = 1$, $\epsilon_2 = \epsilon_3 = \epsilon_4 = 0$, $\epsilon_5 = 1$, $\epsilon_6 = \epsilon_7 = 0$, the matrix is

$$\begin{pmatrix} \binom{1+b-1}{b-1} & \binom{5+b-1}{b-1} & \binom{8+b-1}{b-1} \\ 1 & \binom{4+b-1}{b-1} & \binom{7+b-1}{b-1} \\ 0 & 1 & \binom{3+b-1}{b-1} \end{pmatrix}.$$

When $b = 2$, this is $\begin{pmatrix} 2 & 6 & 9 \\ 1 & 5 & 8 \\ 0 & 1 & 4 \end{pmatrix}$ with determinant 9. Thus the chance of a carry at exactly positions 1 and 5 when adding eight binary numbers is $9/2^8 \doteq .03516$. When $b = 10$, this chance becomes $1,042,470/10^8 \doteq .0104$. This is not so much smaller than in the binary case.

Proof. (Of fact 5) We follow Stanley [67, p. 69]. Let $\alpha_n(S)$ be the number of sequences with descent set contained in S and $\beta_n(S)$ the number of sequences with descent set equal to S . Then

$$\alpha_n(S) = \sum_{T \subseteq S} \beta_n(T), \quad \beta_n(S) = \sum_{T \subseteq S} (-1)^{|S-T|} \alpha_n(T).$$

There is a simple formula for $\alpha_n(S)$: the number of weakly increasing sequences with entries in $\{0, 1, \dots, b-1\}$ of length l is $\binom{l+b-1}{b-1}$ using the usual stars and bars argument (place $b-1$ bars into $l+b-1$ places, put symbol 0 to the left of the first bar, symbol 1 between the first and second bar, and so on, with symbol $b-1$ to the right of the last bar). Then

$$\alpha_n(S) = \binom{s_1+b-1}{b-1} \binom{s_2-s_1+b-1}{b-1} \cdots \binom{n-s_k+b-1}{b-1},$$

because any such sequence of length n is constructed by concatenating nondecreasing sequences of length $s_1, s_2 - s_1, \dots, n - s_k$. Therefore

$$\beta_n(S) = \sum_{1 \leq i_1 < i_2 < \cdots < i_j \leq k} (-1)^{k-j} f(0, i_1) f(i_1, i_2) \cdots f(i_j, k+1)$$

with $f(i, j) = \binom{s_j - s_i + b - 1}{b - 1}$ (with $s_0 = 0$, $s_{k+1} = n$). From Stanley's discussion, $\beta_n(S)$ is the determinant of the $(k+1) \times (k+1)$ matrix with (i, j) entry $f(i, j+1)$, $0 \leq i, j \leq k$. \square

Example 2. From the above development, the chance that the sum of n base b digits has *no* carries is $\binom{n+b-1}{b-1}/b^n$. When $n = 8$, $b = 2$, this is $9/2^8 \doteq .03516$. When $n = 8$, $b = 10$, this is $\binom{17}{9}/10^8 \doteq .00024$. We feel lucky when adding eight numbers with no carries. These calculations show that we are lucky indeed.

3 Determinantal point processes

Let \mathcal{X} be a finite set. A point process on \mathcal{X} is a probability measure P on the $2^{|\mathcal{X}|}$ subsets of \mathcal{X} . For example, if $\mathcal{X} = \{1, 2, \dots, n-1\}$, a point process is specified by recording where the carries occur in adding n base b numbers as in Section 2. One simple way to specify P is via its so-called correlation functions. For $A \subseteq \mathcal{X}$, let

$$\rho(A) = P\{S : S \supseteq A\}.$$

The collection of numbers $\{\rho(A)\}$ uniquely determine P using inclusion/exclusion. A point process is *determinantal* with kernel $K(x, y)$ if

$$\rho(A) = \det (K(x, y))_{x, y \in A}.$$

On the right is the determinant of the $|A| \times |A|$ matrix with (x, y) entry $K(x, y)$ for $x, y \in A$.

Determinantal point processes were introduced by Macchi [50] to model the distribution of fermions. See [17] for a textbook account, [65] for a survey, [41] for probabilistic developments and [48] for many combinatorial examples. We warn the reader that these last references essentially deal with symmetric kernels $K(x, y)$ whereas all of our examples involve non-symmetric kernels.

There has been an explosive development of determinantal point processes over the past few years. Some reasons for this include:

- (a) A raft of natural processes (including all those in the present paper) turn out to be determinantal. These include: the eigenvalues of various ensembles of random matrices (Dyson [25, 26, 27], Mehta [55]); the presence or absence of edges if a random spanning tree is chosen in a graph (Burton–Pemantle [10], Lyons [48]); random tilings and growth models (Johansson [44]); the structure of random partitions chosen from a variety of measures (e.g., the Plancherel measure on the symmetric group S_n) (Borodin–Okounkov–Olshanski [5]); random dimers on bipartite planar graphs (Kenyon [46]); and the zeros of a random analytic function $\sum_{n=0}^{\infty} a_n z^n$ with i.i.d. complex Gaussian coefficients a_n [56]. The list goes on extensively.
- (b) Specifying a kernel may be much easier than specifying a measure on all $2^{|\mathcal{X}|}$ subsets. Further, the correlation functions allow easy computation of quantities of interest. For example, if X_x is 1 or 0 as the random set includes x or not:

$$\begin{aligned} E(X_x) &= K(x, x), \\ \text{Cov}(X_x, X_y) &= \det \begin{pmatrix} K(x, x) & K(x, y) \\ K(y, x) & K(y, y) \end{pmatrix} - K(x, x)K(y, y) \\ &= -K(x, y)K(y, x). \end{aligned}$$

If $K(x, y) = K(y, x)$, then the correlation is ≤ 0 , a distinctive feature of determinantal point processes.

- (c) If the matrix $(K(x, y))_{x, y \in \mathcal{X}}$ has all real eigenvalues $\{\lambda_x\}_{x \in \mathcal{X}}$, many theorems become available for N , the total number of points in a realization of the process. These include:

- N is distributed as a sum $\sum_{x \in \mathcal{X}} Y_x$ with $\{Y_x\}$ independent 0/1 random variables having $P(Y_x = 1) = \lambda_x$.
- Let μ and σ denote the mean and variance of N . These are available in terms of K from (b) above. Then the following refined form of the central limit theorem holds:

$$\left| P \left\{ \frac{N - \mu}{\sigma} \leq x \right\} - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-t^2/2} dt \right| \leq \frac{.80}{\sigma}$$

- The probability density $P(n) = P(N = n)$ is a Polya-frequency function. In particular the density is unimodal. Further, $|\text{mode} - \text{mean}| \leq 1$.
- There is a useful algorithm for simulating from K (here assuming $K(x, y) = K(y, x)$).

Most of these properties follow from the fact that the generating function of N ,

$$E(x^N) = \det(I + (x - 1)K)$$

has all real zeros. See Pitman [57] or Hough, et al. [41] for details.

- (d) If K_n is a sequence of such kernels converging to a kernel K , then, under mild restrictions, the associated point processes converge. See Soshnikov [65] or Johansson and Nordenstam [45] for precise statements and remarkable examples. A simple example is given in Section 5 (Example 11).

A main result of our paper is that the carries and descent processes, and indeed any one-dependent point processes on the integers, is determinantal. The special case of carries is stated here; a proof of the general case is in Section 7.

Theorem 3.1. *The point process of carries and descents P_n given in Section 2 is determinantal with correlation kernel $K(i, j) = k(j - i)$ where*

$$\sum_{m \in \mathbb{Z}} k(m)t^m = \frac{1}{1 - (1 - t)^b}.$$

Example 3. Take $b = 2$. Then $\frac{1}{1 - (1 - t)^2} = \frac{1}{2t(1 - t/2)}$. Replacing t by ct changes $k(m)$ to $k(m)c^m$ and $K(i, j) = k(j - i)$ to $c^{j-i}k(j - i) = c^{-i}K(i, j)c^j$. Conjugating K by a diagonal matrix does not change determinants or the correlation functions. Thus setting $\frac{t}{2} = z$, the generating function becomes

$$\frac{1}{4} \left(\frac{1}{z} + 1 + z + z^2 + \dots \right).$$

When e.g., $n - 1 = 5$, the kernel is

$$\begin{pmatrix} \frac{1}{4} \\ \frac{1}{4} \end{pmatrix} \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix}. \quad (3.1)$$

From this, $\rho(A) = \begin{cases} 0 & \text{if } A \text{ has two consecutive entries} \\ (\frac{1}{4})^{|A|} & \text{otherwise} \end{cases}$. This is a manifestation of the one-dependence together with the fact that (for $b = 2$), two consecutive carries are impossible. It follows that the *possible* configurations are all binary strings of length $n - 1$ with no two consecutive 1's. This is a standard coding of the Fibonacci number F_{n+1} (if 1, 1, 2, 3, 5 are the first five Fibonacci numbers). For example, when $n = 4$, the possible configurations are 000, 001, 010, 100, 101.

The eigenvalues of any finite matrix of the form (3.1) are real. This follows from the fact that these matrices are totally positive, see [28], and the well known fact that the eigenvalues of a totally positive matrix are real (and nonnegative), see e.g. Corollary 6.6 of [1].

The classification of [28] also implies that the correlation kernels for $b \geq 3$ are not totally positive (for large enough matrix size), and we do not know if their eigenvalues are real.

Example 4. Take $b = 3$. A straightforward expansion leads to

$$\begin{aligned} \sum_{m=-\infty}^{\infty} k(m)t^m &= \frac{1}{1 - (1-t)^3} = \frac{1}{3t \left(1 - t + \frac{t^2}{3}\right)} \\ &= \frac{\left(1 + t + \frac{2t^2}{3} + \frac{t^3}{3} + \frac{t^4}{9}\right)}{3t \left(1 + \frac{t^6}{27}\right)}. \end{aligned}$$

Thus

- $k(n) = 0$ for $n < -1$,
- $k(-1) = \frac{1}{3}$,
- $k(n) = 0$ for $n = 4 + 6j$, $0 \leq j < \infty$. For other $n \geq 2$,
- $k(n) = (-1)^{\lfloor \frac{n+1}{6} \rfloor} \left(\frac{1}{3}\right)^{\lfloor \frac{n+3}{2} \rfloor} 2^{\delta(n)}$, where $\delta(n) = 1$ if $n = 1 \pmod 6$ and 0 else.

For example, the first few values are:

m	-1	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
$k(m)$	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{2}{9}$	$\frac{1}{9}$	$\frac{1}{27}$	0	$-\frac{1}{3^4}$	$-\frac{1}{3^4}$	$-\frac{2}{3^5}$	$-\frac{1}{3^5}$	$-\frac{1}{3^6}$	0	$\frac{1}{3^7}$	$\frac{1}{3^7}$	$\frac{2}{3^8}$	$\frac{1}{3^8}$	$\frac{1}{3^9}$	0

It is amusing to see that the two-variable kernel $K(x, y) = k(y - x)$ reproduces the correlation functions found for this problem in Section 2 (Fact 4). Thus for $n = 7$,

$$(K(x, y))_{i,j=1}^6 = \begin{bmatrix} 1/3 & 2/9 & 1/9 & 1/27 & 0 & -1/81 \\ 1/3 & 1/3 & 2/9 & 1/9 & 1/27 & 0 \\ 0 & 1/3 & 1/3 & 2/9 & 1/9 & 1/27 \\ 0 & 0 & 1/3 & 1/3 & 2/9 & 1/9 \\ 0 & 0 & 0 & 1/3 & 1/3 & 2/9 \\ 0 & 0 & 0 & 0 & 1/3 & 1/3 \end{bmatrix}$$

4 One-dependent processes

Let $\{X_i\}_{i=0}^N$ be a binary stochastic process. Here $0 < N \leq \infty$. The process is *one-dependent* if $\{X_j\}_{j=0}^{i-1}$, $\{X_j\}_{j=i+1}^N$ are independent for all i . A natural example: let $\{Y_i\}_{i=0}^{N+1}$ be independent uniform random variables on $[0, 1]$. Let $h : [0, 1]^2 \rightarrow \{0, 1\}$ be a measurable function. Then $X_i = h(Y_i, Y_{i+1})$, $0 \leq i \leq N$, is a one-dependent process called a *two-block factor*. Much of the literature on one-dependence is for stationary

processes. However, if Y_i are independent but not identically distributed or $h = h_i$ depends on i , the associated X_i is still one-dependent. Our main results in Section 7 are for general one-dependent processes. We first describe some results for the stationary case.

Stationary one-dependent processes

A book-length review of this subject by de Valk [73] collects together many of the results, so we will be brief, focusing on later research. One of the first problems studied is the question of whether all one-dependent processes are two-block factors. Counterexamples were found [73] though it appears that most “natural” one-dependent processes are two-block factors. For example, let $f \in L^2(\pi)$ be a function on the unit circle. Let $K(i, j) = \hat{f}(i - j)$, $i, j \in \mathbb{Z}$, \hat{f} the Fourier transform. Macchi [50] and Soshnikov [65] show that $K(i, j)$ is the kernel for a determinantal point process on \mathbb{Z} . Lyons and Steif [49] and Shirai and Takahashi [62, 63] develop remarkable properties of the associated point process. If $f(\theta) = ae^{-i\theta} + b + ce^{i\theta}$, the point process is one-dependent and [9] shows it is a two-block factor. See also [13].

A stationary one-dependent process is determined by the numbers $a_i = P(X_1 = X_2 = \dots = X_{i-1} = 1)$, $a_1 = 1$. Indeed, if $f = f_1, \dots, f_{i-1}; e = e_{i+1}, \dots, e_{n-1}$ are binary strings, $P(f, 0, e) + P(f, 1, e) = P(f)P(e)$. From this $P(f, 0, e) = P(f)P(e) - P(f, 1, e)$. By similar reductions, the probability of any pattern of occurrences can be reduced to a polynomial in the a_i . For example, $P(0, 0, 0) = 1 - 3a_2 + a_2^2 + 2a_3 - a_4$. There is a remarkable, simple formula for this polynomial as a minor of a Toeplitz matrix. We learned this from [58, Chap. 7]. A generalization to the non-stationary case is in Section 7.

Theorem 4.1. *For a stationary one-dependent process, let $a_i = P(X_1 = X_2 = \dots = X_{i-1} = 1)$, $a_1 = 1$. Let t_1, \dots, t_{n-1} be a binary string with j zeros at positions $S = \{s_1 < \dots < s_k\} \subseteq [n - 1]$. Then*

$$P(t_1, \dots, t_{n-1}) = \det (a_{s_{j+1} - s_i})_{i,j=0}^k.$$

Here the determinant is of a $k+1$ by $k+1$ matrix, and one sets $s_0 = 0, s_{k+1} = n, a_0 = 1$, and $a_i = 0$ for $i < 0$.

Proof. Theorem 4.1 follows from Proposition 2.2 and Theorem 4.2 in [58, Chap. 7], after elementary manipulations. \square

Remark: It follows from the discussion on [58, p. 140] that if all of the determinants (for all n and subsets S) in Theorem 4.1 are non-negative, then a stationary, one-dependent process with these a_i values exists.

Example 5. For any n , the sequence with $n - 1$ ones has $k = 0$ zeros. The relevant matrix is then 1 by 1 with entry a_n , giving this probability.

Example 6. To compute $P(0, 0, 0)$ one applies the theorem with $n = 4$ and $S = \{1, 2, 3\}$. It follows that

$$P(0, 0, 0) = \det \begin{bmatrix} 1 & a_2 & a_3 & a_4 \\ 1 & 1 & a_2 & a_3 \\ 0 & 1 & 1 & a_2 \\ 0 & 0 & 1 & 1 \end{bmatrix} = 1 - 3a_2 + a_2^2 + 2a_3 - a_4,$$

as above.

Our next result expresses the determinant in Theorem 4.1 as a skew-Schur function of ribbon (also called rim-hook) type. Background on skew-Schur functions can be found in [51, Sec. 1.5]. For further references and recent work, one can consult [3] or [47]. We will use the elementary symmetric functions e_r in countably many variables defined as $e_0 = 1$ and

$$e_r = \sum_{1 \leq i_1 < i_2 < \dots < i_r} x_{i_1} x_{i_2} \dots x_{i_r}.$$

From [51, p. 20] these are algebraically independent, so there is a homomorphism of the ring of symmetric functions to \mathbb{R} which sends each e_i to a_i from Theorem 4.1.

Theorem 4.2. *With notation as in Theorem 4.1, let λ and μ be the partitions defined by*

$$\lambda_i = n - s_{i-1} - k + i - 1, \quad \mu_i = n - s_i - k + i - 1, \quad 1 \leq i \leq k + 1.$$

Let λ', μ' denote the transpose partitions of λ and μ and let $s_{\lambda'/\mu'}$ denote the corresponding skew-Schur function, obtained by specializing the elementary symmetric functions e_i to equal a_i . Then

$$P(t_1, \dots, t_{n-1}) = s_{\lambda'/\mu'}.$$

Proof. First note that the λ_i, μ_i defined above are all non-negative. From Macdonald [51, p. 71],

$$s_{\lambda'/\mu'} = \det(e_{\lambda_i - \mu_j - i + j})_{i,j=1}^{k+1}, \quad (4.1)$$

and e_r is the r th elementary symmetric function. When each e_r is specialized to equal a_r , the quantity $s_{\lambda'/\mu'}$ becomes

$$\det(a_{\lambda_i - \mu_j - i + j})_{i,j=1}^{k+1} = \det(a_{s_j - s_{i-1}})_{i,j=1}^{k+1},$$

as desired. □

One reason why Theorem 4.2 is interesting is that there are non-obvious equalities between ribbon skew-Schur functions. The paper [3] characterizes when two ribbon skew-Schur functions are equal; analogous results for more general skew-Schur functions are in [60]. In particular, combining the results of [3] with Theorem 4.2 one immediately obtains the fact that a stationary one-dependent process is invariant under time reversal, i.e.

$$P(t_1, \dots, t_n) = P(t_n, \dots, t_1)$$

for all t_i . For another proof of invariance under time reversal, see [58, p. 139].

The determinantal formulae of Theorems 4.1 and 4.2 suggest that a determinantal point process is lurking nearby. This is indeed the case. In Section 7 (see Corollary 7.3) we prove the following.

Corollary 4.3. *A stationary one-dependent process as in Theorem 4.1 is determinantal with kernel $K(x, y) = k(y - x)$ with*

$$\sum_{n \in \mathbb{Z}} k(n) z^n = -1 / \sum_{j=1}^{\infty} a_j z^j.$$

A remarkable development, connecting stationary one-dependent processes to algebras appears in [58, Chap. 7]. They consider a graded algebra $A_0 \oplus A_1 \oplus A_2 + \dots$ with $A_0 = k$, a ground field, and $A_i A_j \subseteq A_{i+j}$. For example, the space $k[x]$ of polynomials in one variable has $A_0 = k$, and A_i spanned by x^i . They assume that each A_i is a finite dimensional vector space of dimension $\dim(A_i)$. The algebra is *quadratic* if $A_0 = k$, the algebra is generated by elements of A_1 , and the relations defining the algebra are in A_2 . For example, the commutative polynomial ring $k[x_1, \dots, x_n]$ is generated by x_1, \dots, x_n and the quadratic relations $x_i x_j - x_j x_i = 0$. Note that algebras need not be commutative.

A technical growth condition on the $\dim(A_i)$ which we will not explain here yields the Koszul algebras. These include *many* natural algebras occurring in mathematics (see [34] for a survey). As a simple example, consider the commutative polynomial ring generated by x_1, x_2, \dots, x_n with the additional relations $x_i x_j = 0$ for $(i, j) \in E$, with E the edge set of an undirected graph on $\{1, 2, \dots, n\}$ (loops allowed). This is Koszul [58, Chap. 2, Cor. 4.3]. Polishchuk and Positselski [58, Chap. 7, Cor. 4.3] prove the following remarkable result.

Theorem 4.4. *To every Koszul algebra A one can assign a stationary, one-dependent process via*

$$P(X_1 = X_2 = \dots = X_{i-1} = 1) = \frac{\dim(A_i)}{\dim(A_1)^i} \quad \text{for } i = 1, 2, \dots$$

Part of the reason that Theorem 4.4 is substantive is that the quantities $P(X_1 = X_2 = \dots = X_{i-1} = 1)$ can not assume arbitrary values; the determinants in Theorem 4.1 must be non-negative.

Example 7. Consider the commutative polynomial algebra generated by x_1, x_2, \dots, x_b and the additional relations $x_i^2 = 0$, $1 \leq i \leq b$. The degree i part A_i is spanned by square free monomials and has $\dim(A_i) = \binom{b}{i}$, $0 \leq i < \infty$. From Fact 2 of Section 2, we see that the associated one-dependent process is precisely the carries process of mod b addition from our introduction. Note that since only the $\dim(A_i)$ matter, the algebra in this example can be replaced by the exterior algebra generated by x_1, x_2, \dots, x_b and the relations $x_i x_j = -x_j x_i$ for all i, j .

Example 8. Consider the commutative polynomial algebra generated by x_1, x_2, \dots, x_b . The degree i part A_i is spanned by monomials of degree i and has $\dim(A_i) = \binom{b+i-1}{i}$. From Example 2 of Section 2, the associated one-dependent process is precisely the complement of the carries process of mod b addition. More generally, in Section 7 we show that the particle-hole involution of a (perhaps non-stationary) one-dependent process with respect to any subset is one-dependent.

Example 9. A PBW algebra is an algebra of the form

$$k[x_1, \dots, x_n] / \langle x_i x_j : (i, j) \in S \subset [1, n]^2 \rangle,$$

where the variables x_1, \dots, x_n do not commute. Here it is useful to think of S as a directed graph with vertex set $\{1, \dots, n\}$, loops allowed. From [58, p. 84], PBW algebras are Koszul. As noted in [58, Chap. 7, Prop. 5.1], the one-dependent process

associated to a PBW algebra can be described as follows: pick U_1, U_2, \dots i.i.d. in $\{1, \dots, n\}$ and let $X_i = h(U_i, U_{i+1})$, where $h(i, j) = 0$ if and only if $(i, j) \in S$. Indeed, the dimensions satisfy $\dim(A_0) = 1, \dim(A_1) = n, \dim(A_i) = \text{number of paths of length } i - 1 \text{ in the complement of } S$, and so the chance of $i - 1$ consecutive 1's in the point process is equal to $\dim(A_i) / \dim(A_1)^i$.

The above argument shows that PBW algebras give rise to two-block-factor processes. In fact they are dense in the set of all two-block factors [58, Chap. 7, Prop. 5.2]. For a generalization of these PBW processes, as well as a simple argument that they are determinantal, see Remark 5 after Theorem 5.3.

Example 10. Consider $2n$ points x_1, x_2, \dots, x_{2n} in general position in projective space \mathbb{P}^n . The coordinate ring of this projective variety is known to be Koszul [58, p. 42] with $\dim(A_0) = 1, \dim(A_1) = n + 1, \dim(A_i) = 2n$ for $i \geq 2$. Thus $P(X_1 = X_2 = \dots = X_{i-1} = 1) = 2n / (n + 1)^i$ defines a one-dependent process. When $n = 1$, this is fair coin tossing. When $n = 3$, it is the carries process for multiplying quaternions introduced in Section 6.

For general n , these one-dependent processes can be described as follows (and illustrates what one might consider to be a “probabilistic” description). Let $\{1, 2, \dots, n\}$ be ordered cyclically and let $*$ be another symbol. Choose U_i i.i.d. in $\{*, 1, \dots, n\}$. Let $X_i = h(U_i, U_{i+1})$ where $h(*, x) = 0$ for all x , $h(x, *) = 1 = h(i, i + 1)$ for $x \neq *$, and $h(i, j) = 0$ otherwise. To see that this works, note that $P(X_1 = 1) = 2n / (n + 1)^2$, since one has to choose x different from $*$ first and then $*$ or $x + 1$ next. Similarly, for the chance of $i - 1$ 1's in a row, the first choice can be anything but $*$, then the next $i - 2$ choices are determined and the last can be one of two.

We are certain that natural Koszul algebras will lead to natural point processes. We note further that [58, Sec. 7.6] shows how natural operations on algebras preserve the Koszul property. These include the operations of union and complements that we work with in Section 7.

5 Descents in permutations

A permutation $\sigma \in S_n$, the symmetric group, has a descent at i if $\sigma(i) > \sigma(i + 1)$. The set of such i forms the descent set $D(\sigma)$. If the base b in previous sections is large and n stays fixed, a string of n digits will have no repeated values and the descent theory of Section 2 and Section 3 above becomes descent theory for random permutations. This is a venerable subject. Stanley [66, 68] reviews the basics. Some modern highlights are Solomon's descent algebra [64], the connections with the free Lie algebra [36, 61], Gessel's theory of enumerating permutations by descents and cycle structure [22, 37], quasi-symmetric functions and the theory of riffle shuffling [35, 70]. Any Coxeter group has its own descent theory [64]. There is some indication that arithmetic carries can be carried over [20].

This section introduces three examples where the descent set can be shown to be a one-dependent determinantal point process: uniform choice of permutations, non-uniform choice from the Mallows model, and independent trials. Then it shows that the closely related notion of the connectivity set of a permutation also yields a determinantal process. Finally, connections with binomial posets are mentioned.

Example 11. The descent set of a uniformly chosen permutation. Consider the formula of Section 2 Fact 5 for the chance that a random b -ary string of length n has descents exactly at $S \subseteq [n-1]$. Passing to the limit, as $b \nearrow \infty$ using $\binom{a+b-1}{b-1} \sim \frac{b^a}{a!}$ gives a classical formula for the chance that a uniformly chosen $\sigma \in S_n$ has descents exactly at S :

$$P_n(D(\sigma) = S) = \det [1/(s_{j+1} - s_i)!]. \quad (5.1)$$

For $S = 1 \leq s_1 < s_2 < \dots < s_k \leq n-1$, the determinant is of a $(k+1) \times (k+1)$ matrix with (i, j) entry $1/(s_{j+1} - s_i)!$, $(i, j) \in [0, k] \times [0, k]$ and $s_0 = 0, s_{k+1} = n$. This formula is originally due to MacMahon [52]. See Stanley [67, p. 69] or Gessel and Viennot [38] for modern proofs.

Equation (5.1) allows us to see that the descent set of a uniformly random permutation, treated as a point process, is determinantal. The proof follows from (5.1) and Corollary 7.6 in Section 7.

Theorem 5.1. *There exists a stationary point process on \mathbb{Z} , call it P , such that its restriction to any interval of length $n-1$ coincides with P_n of (5.1). The process P is one-dependent and determinantal. Its correlation kernel $K(x, y) = k(y-x)$ with*

$$\sum_{m \in \mathbb{Z}} k(m)z^m = \frac{1}{1-ez}. \quad (5.2)$$

Remarks.

1. Theorem 5.1 can also be seen by passing to the limit in Theorem 3.1. Replacing t in the generating function $1/(1-(1-t)^b)$ by $-z/b$ and letting $b \nearrow \infty$ gives (5.2).
2. The Bernoulli numbers B_n are defined by $z/(e^z-1) = \sum_{n=0}^{\infty} B_n z^n/n!$. These are very well-studied. It is known that $B_{2i+1} = 0$ for $i \geq 1$, $B_0 = 1, B_1 = -\frac{1}{2}, B_2 = \frac{1}{6}, B_4 = -\frac{1}{30}, B_6 = \frac{1}{42}, \dots$. We see that $k(m) = -B_{m+1}/(m+1)!$.
3. From these calculations, the kernel K is

$$K = \begin{bmatrix} \frac{1}{2} & -\frac{1}{12} & 0 & \frac{1}{720} & \dots \\ -1 & \frac{1}{2} & -\frac{1}{12} & 0 & \ddots \\ & -1 & \ddots & \ddots & \ddots \\ & & \ddots & \ddots & \ddots \\ & & & \ddots & \ddots \end{bmatrix}.$$

4. The correlation functions of P (and P_n) can be computed explicitly by passage to the limit from Fact 5 of Section 2. Let $A = A_1 \cup A_2 \cup \dots \cup A_k$ be a decomposition of the finite set $A \subseteq \mathbb{Z}$ into disjoint blocks of adjacent integers. If $|A_i| = a_i$,

$$\rho(A) = \prod_{i=1}^k \frac{1}{(a_i+1)!}.$$

5. For any $n = 1, 2, \dots$, let $K_n(i, j) = k(j - i)$ be the $n \times n$ top left corner block of K . Let $d(\pi)$ denote the number of descents of π and let

$$A_{n+1} = \sum_{\pi \in S_{n+1}} x^{d(\pi)}$$

be the $(n+1)$ st Eulerian polynomial (see e.g., [67, p. 22]). Then item (c) preceding Theorem 3.1 yields that

$$A_{n+1}(x) = \det(I + (x - 1)K_n) (n + 1)!$$

It is known [14, p. 292] that $A_{n+1}(x)$ has all real zeros $\alpha_1, \alpha_2, \dots, \alpha_n$. From the development in Section 3, N_{n+1} , the number of descents in a random permutation from S_{n+1} , is the sum of n independent Bernoulli random variables with success rates $(1 - \alpha_j)^{-1}$, $1 \leq j \leq n$. One easily computes

$$\begin{aligned} E(N_{n+1}) &= \text{tr}(K_n) = \frac{n}{2}, & \text{Var}(N_{n+1}) &= \text{tr}(K_n - K_n^2) \\ & & &= \frac{n}{2} - \left(\frac{5n}{12} - \frac{1}{6}\right) = \frac{n+2}{12}. \end{aligned}$$

It follows by Harper's method ([39],[57]) that one has a central limit theorem:

$$\frac{N_{n+1} - \frac{n}{2}}{\sqrt{\frac{n+2}{12}}} \implies \mathcal{N}(0, 1).$$

6. Chebikin [11] studied the ‘‘alternating descent set’’ $A(\sigma)$ of a random permutation, defined as the set of positions at which the permutation has an *alternating descent*, that is an ordinary descent if i is odd, or an ascent if i is even. Combining Lemma 2.3.1 of [11] with the argument used to prove (5.1), one obtains that for $|S| = k$,

$$P_n(A(\sigma) = S) = \det[E_{s_{j+1}-s_i}/(s_{j+1} - s_i)!]_{i,j=0}^k,$$

where E_n is the n th Euler number defined by $\sum_{n \geq 0} E_n z^n / n! = \tan(z) + \sec(z)$, and $s_0 = 0, s_{k+1} = n$. Applying our Theorem 7.5 in Section 7, it follows that that P_n is obtained by restricting to any interval of length $n - 1$ the stationary, one-dependent, determinantal process with correlation kernel $K(x, y) = k(y - x)$ with

$$\sum_{m \in \mathbb{Z}} k(m)z^m = \frac{1}{1 - (\tan(z) + \sec(z))^{-1}}.$$

7. Modern combinatorics suggests a host of potential generalizations of Theorem 5.1. Let P be a partial order on $[n]$. A linear extension of P is a permutation $\sigma \in S_n$ such that if i is less than j according to P , then $\sigma(i) < \sigma(j)$. Let $\mathcal{L}(P)$ denote the set of linear extensions of P . For each $\sigma \in \mathcal{L}(P)$, let

$$\begin{aligned} D(\sigma) &= \{i : \sigma(i) > \sigma(i + 1)\}, & \text{the descent set of } \sigma, \\ \wedge(\sigma) &= \{i : \sigma(i - 1) < \sigma(i) > \sigma(i + 1)\}, & \text{the peak set of } \sigma. \end{aligned}$$

Choosing σ uniformly in $\mathcal{L}(P)$ gives two point processes. This section has focused on ordinary descents, that is, P is the trivial poset with no restrictions. There are many indications that ‘‘natural’’ posets will give rise to determinantal point processes. For background and first results, see Brenti [8] or Stembridge [72].

8. Finally, we mention that it would be interesting to find analogs of Theorem 5.1 for other Coxeter groups. For the hyperoctahedral group B_n consisting of the $2^n n!$ signed permutations, descents can be defined using the linear ordering

$$1 < 2 < \cdots < n < -n < \cdots < -2 < -1.$$

Say that

- (a) σ has a descent at position i ($1 \leq i \leq n-1$) if $\sigma(i) > \sigma(i+1)$.
- (b) σ has a descent at position n if $\sigma(n) < 0$.

For example, $-4 -1 3 2 5 \in B_3$ has descent set $\{2, 3, 5\}$. Reiner [59] shows that the chance that a random element of B_n has descent set $\{s_1, \dots, s_r\} \subseteq \{1, 2, \dots, n\}$ is $\det(a(i, j))_{i, j=0}^r$ where

$$a(i, j) = \begin{cases} \frac{1}{(s_{j+1} - s_i)!} & \text{if } 0 \leq j \leq r-1 \\ \frac{1}{2^{n-s_i} (n-s_i)!} & \text{if } j = r \end{cases}$$

Here $s_0 = 0$. Theorem 7.5 in Section 7 shows that the resulting process is determinantal (but not stationary).

Example 12. Descents from non-uniform distributions on permutations. In a variety of applications, *non-uniform* distributions are used on permutations. A widely used model is

$$P_\theta(\sigma) = Z^{-1} \theta^{d(\sigma, \sigma_0)}, \quad \sigma \in S_n. \quad (5.3)$$

Here, $0 < \theta \leq 1$ is a fixed concentration parameter, $\sigma_0 \in S_n$ is a fixed location parameter, $d(\sigma, \sigma_0)$ is a metric on S_n and $Z = \sum_{\sigma} \theta^{d(\sigma, \sigma_0)}$ is the normalizing constant. These are called “Mallows models through the metric d .” When $\theta = 1$, P_1 is the uniform distribution. For $0 < \theta < 1$, P_θ is largest at σ_0 and falls off as σ moves away from σ_0 . See [15], [23] or [54] for background and references.

Perhaps the most widely used metric is

$$\begin{aligned} d(\sigma, \sigma_0) &= \text{minimum number of pairwise adjacent transpositions} \\ &\quad \text{to bring } \sigma \text{ to } \sigma_0, \\ &= I(\sigma \sigma_0^{-1}), \text{ the number of inversions in } \sigma \sigma_0^{-1}. \end{aligned} \quad (5.4)$$

This is called “Kendall’s tau” in the statistics literature. References, extensions and properties are in [19, Sect. 4], [31], [53]. We mention that the normalizing constant of the Mallows model through Kendall’s tau is given by $Z = \prod_{i=1}^n \frac{\theta^i - 1}{\theta - 1}$ [67, p. 21].

With θ, σ_0 and d fixed, one may ask any of the usual questions of applied probability and enumerative combinatorics: “Picking a permutation randomly from $P_\theta(\cdot)$, what is the distribution of the cycle structure, longest increasing subsequence, . . . ?” For general metrics, these are open research problems. Even the algorithmic task of sampling from P_θ leads to difficult problems. See [21] or [19].

For Mallows model P_θ through Kendall’s tau (5.4) and $\sigma_0 = \text{id}$, we show that the descent set of σ , $D(\sigma) = \{i : \sigma(i+1) < \sigma(i)\}$ forms a determinantal point process with simple properties. In what follows we use the q -analog notation that $i_q = \frac{q^i - 1}{q - 1}$ and $n!_q = \prod_{i=1}^n i_q$, and take $\theta = q$ as is conventional in combinatorial work in the subject.

Proposition 5.2. Let P_q ($0 < q < 1$) be Mallows model through Kendall's tau (5.3), (5.4) on S_n .

(a) The chance that a random permutation chosen from P_θ has descent set $s_1 < s_2 < \dots < s_k$ is

$$\det \left[\frac{1}{(s_{j+1} - s_i)!_q} \right]_{i,j=0}^k,$$

with $s_0 = 0$ and $s_{k+1} = n$.

(b) The point process associated to $D(\sigma)$ is stationary, one-dependent, and determinantal with kernel $K(x, y) = k(y - x)$, where

$$\sum_{m \in \mathbb{Z}} k(m) z^m = \frac{1}{1 - (\sum_{m=0}^{\infty} z^m / m!_q)^{-1}}.$$

(c) The chance of finding k descents in a row is $q \binom{k+1}{2} / (k+1)!_q$. In particular, the chance of a descent at any given location is $q/(q+1)$. The number of descents has mean $\mu(n, q) = \frac{(n-1)q}{q+1}$ and variance $q \left[\frac{(q^2 - q + 1)n - q^2 + 3q - 1}{(q^2 + q + 1)(q+1)^2} \right]$. Normalized by its mean and variance, the number of descents has a limiting standard normal distribution.

Proof. Part (a) of Proposition 5.2 follows from Stanley [67, Ex. 2.2.5]. Part (b) follows from Corollary 7.6 in Section 7 and elementary calculations. To compute the chance of k consecutive descents, note by stationarity that this is the chance of the permutation $\sigma(i) = k+2-i$ in S_{k+1} under the P_q measure on S_{k+1} . This probability is $q \binom{k+1}{2} / (k+1)!_q$, as needed. One calculates from part (b) that

$$k(-1) = 1, \quad k(0) = \frac{q}{q+1}, \quad k(1) = \frac{q^2}{(q+1)^2(q^2 + q + 1)}.$$

Recall from the remarks preceding Theorem 3.1 that

$$E(x^N) = \det(I + (x-1)K),$$

where N is the total number of particles. Thus

$$E(|D(\sigma)|) = \text{tr}(K_{n-1}) = (n-1)k(0),$$

$$\text{Var}(|D(\sigma)|) = \text{tr}(K_{n-1} - K_{n-1}^2) = (n-1)k(0) - (n-1)k(0)^2 - 2(n-2)k(1)k(-1),$$

and the proof of part (c) is complete. \square

Remarks.

1. From part (c), the correlation functions are given explicitly, just as in remark 4 following Theorem 5.1.
2. It would be interesting to have a direct combinatorial proof of the one-dependence in Proposition 5.2.

Example 13. Descents for independent trials The carries in the carries process have an equivalent description in terms of descents in an independent and uniform sequence. In this example we generalize to descents in a sequence Y_1, Y_2, \dots, Y_n with Y_i independent and identically distributed with $P(Y_1 = j) = p_j$. Here $0 \leq j \leq b-1$ to match previous sections (and of course $0 \leq p_j \leq 1, \sum p_j = 1$), but in fact $b = \infty$ can be allowed. The descents are related to the order statistics for discrete random variables [2, 29] but we have not seen this problem previously treated in the probability or statistics literature. Of course, if Y_i are independent and identically distributed from a continuous distribution, the descent theory is the same as the descent theory for a random permutation.

It is useful to have the complete homogeneous symmetric functions

$$h_k(x_1, x_2, \dots, x_m) = \sum_{1 \leq i_1 \leq \dots \leq i_k \leq m} x_{i_1} x_{i_2} \dots x_{i_k}.$$

Theorem 5.3. *If Y_1, Y_2, \dots, Y_n are independent with common probability $P(Y_1 = j) = p_j$, $0 \leq j \leq b-1$, the descents in Y_1, Y_2, \dots, Y_n form a stationary one-dependent point process on $[n-1]$ with*

$$P_n(S) = \det (h_{s_{j+1}-s_i})_{0 \leq i, j \leq k} \quad \text{with } s_0 = 0, s_{k+1} = n,$$

for $S = \{s_1 < s_2 < \dots < s_k\}$. The h_j are evaluated at p_0, \dots, p_{b-1} . Moreover, this process is determinantal with $K(i, j) = k(j-i)$ and $\sum_{l=-\infty}^{\infty} k(l)z^l = \frac{1}{1 - \prod_{i=0}^{b-1} (1-p_i z)}$.

Proof. Let $\alpha_n(S) = P_n(\text{des}(Y_1, Y_2, \dots, Y_n) \subseteq S) = h_{s_1} h_{s_2-s_1} \dots h_{n-s_k}$. The determinant formula follows from the argument for Fact 5 in Section 2. Now, Corollary 7.6 implies that $\hat{k}(z) = \frac{1}{1-\hat{e}(z)^{-1}}$, with $\hat{e}(z) = \sum_{l=0}^{\infty} h_l z^l = \prod_{i=0}^{b-1} (1-p_i z)^{-1}$ from e.g., Macdonald [51, p. 21]. \square

Remarks.

1. Setting $p_i = 1/b$, $\hat{k}(z) = \frac{1}{1-(1-z/b)^b}$. This is equivalent to our result in Theorem 3.1, by changing variable z to $z/b \rightarrow t$, which doesn't change correlations.
2. From the theorem, the associated point process is one-dependent. If $X_i = 1$ or 0 as there is a descent at i or not, then clearly

$$P(X_1 = X_2 = \dots = X_i = 1) = e_{i+1}(p_0, \dots, p_{b-1}),$$

where $e_r(x_1, \dots, x_m) = \sum_{1 \leq i_1 < \dots < i_r \leq m} x_{i_1} x_{i_2} \dots x_{i_m}$ is the r th elementary symmetric function. For example,

$$P(X_i = 1) = \frac{1}{2} - \frac{1}{2} \sum_i p_i^2,$$

$$P(X_i = X_{i+1} = 1) = \frac{1}{6} - \frac{1}{2} \sum_i p_i^2 + \frac{1}{3} \sum_i p_i^3.$$

This allows a straightforward formulation (and proof) of the central limit theorem for the number of descents. As in Section 2, simple expressions for the correlation functions are available in terms of $P(X_1 = X_2 = \dots = X_i = 1)$.

3. As we show in Corollary 7.7, one can express $P_n(S)$ as a skew-Schur function of ribbon type. For special choices of the p_i 's, this symmetric function interpretation may facilitate computation. For example, [51, Sect. 1.5] gives $s_{\lambda/\mu} = \sum_{\nu} c_{\mu\nu}^{\lambda} s_{\nu}$ with $c_{\mu\nu}^{\lambda}$ the Littlewood–Richardson coefficients and s_{ν} the Schur function. If $p_i = zq^i$ ($i = 0, 1, \dots, b-1$, z a normalizing constant), then s_{ν} is non-0 only if ν has length $\leq b$, in which case

$$s_{\nu} = z^{|\nu|} q^{n(\nu)} \prod_{x \in \nu} \frac{1 - q^{b+c(x)}}{1 - q^{h(x)}};$$

see Macdonald [51, p. 44] for an explanation of notation.

4. Coming back to carries, if a column of numbers is chosen independently on $\{0, 1, \dots, b-1\}$ from common distribution $\{p_i\}_{i=0}^{b-1}$, the remainder column after addition evolves as a Markov chain with starting distribution $\{p_i\}$ and transition matrix $P(i, j) = p_{j-i}$ (indices mod b). This is a circulant with known eigenvalues and eigenvectors. We do not know the distribution of carries.
5. There is a sweeping generalization of this example which leads to a large collection of determinantal point processes. Let R be an arbitrary subset of $[N] \times [N]$. Fix a probability distribution $\theta_1, \theta_2, \dots, \theta_N$ on $[N]$. A point process P_n on $[n-1]$ arises as follows: pick Y_1, Y_2, \dots, Y_n independently from $\{\theta_i\}$. Let $X_i = \begin{cases} 1 & \text{if } (Y_i, Y_{i+1}) \notin R \\ 0 & \text{otherwise} \end{cases}$, $1 \leq i \leq n-1$. Let $S = \{i : X_i = 1\}$. Using essentially the arguments above, it is proved in Brenti [8, Th. 7.32] that

$$P_n(S) = \det \left[h_{s_{j+1}-s_i}^R \right]_{i,j=0}^k \quad \text{for } S = \{s_1 < \dots < s_k\}, s_0 = 0, s_{k+1} = n,$$

where $h_j^R = h_j^R(\theta_1, \dots, \theta_N) := P_j(\emptyset)$. Here as usual, $h_0^R = 1, h_j^R = 0$ for $j < 0$. This falls into the domain of Theorem 7.5 and its corollary. It follows that P_n is determinantal with kernel $K(i, j) = k(j-i)$, $\sum_{l \in \mathbb{Z}} k(l)z^l = \frac{1}{(1 - (\sum_{j=0}^{\infty} h_j^R z^j)^{-1})}$.

Our examples are the special case $R(i, j) = \begin{cases} 1 & \text{if } i \leq j \\ 0 & \text{otherwise} \end{cases}$. We are certain that there are many other interesting cases.

Example 14. Connectivity set of a permutation.

Stanley [71] studied a “dual” notion to the descent set of a permutation, which he called the connectivity set. For $\sigma \in S_n$, the connectivity set $C(\sigma)$ is defined as the set of i , $1 \leq i \leq n-1$, such that $\sigma \in S_{1, \dots, i} \times S_{i+1, \dots, n}$. For example, the permutation $\sigma = 3 \ 1 \ 2 \ 5 \ 4 \ 6$ satisfies $C(\sigma) = \{3, 5\}$. The connectivity set also arises in the analysis of quicksort, where it is called the set of splitters of a permutation [74, Sec. 2.2].

We will prove that the connectivity set of a random element of S_n is a determinantal point process and determine its correlation kernel. For this the following two facts are helpful:

1. A permutation $\sigma \in S_n$ is called *indecomposable* or *connected* if $C(\sigma) = \emptyset$. Comtet [14, Exer. VII.16] shows that the number $f(n)$ of connected permutations in S_n satisfies

$$\sum_{n \geq 1} f(n)x^n = 1 - \frac{1}{\sum_{n \geq 0} n!x^n}. \quad (5.5)$$

(An asymptotic expansion of $f(n)$ is given in [14, Exer. VII.17]).

2. [71, Prop. 1.1] Letting $S = \{s_1 < s_2 < \dots < s_k\}$ be a subset of $\{1, 2, \dots, n-1\}$,

$$|\{\sigma \in S_n : S \subseteq C(\sigma)\}| = s_1!(s_2 - s_1)! \cdots (s_k - s_{k-1})!(n - s_k)!. \quad (5.6)$$

The following is our main result.

Theorem 5.4. *Let $C(\sigma)$ denote the connectivity set of a permutation σ chosen uniformly at random from S_n . The point process corresponding to $C(\sigma) \cup \{0, n\}$ is determinantal, with state space $\{0, \dots, n\}$ and correlation kernel $K(x, y)$ satisfying*

$$\begin{aligned} K(0, y) &= 1 && \text{all } y \\ K(n, y) &= \delta_{n,y} && \text{all } y \\ K(x, y) &= \frac{1}{\binom{n}{x}} && 0, n \neq x \leq y \\ K(x, y) &= \frac{1}{\binom{n}{x}} - \frac{1}{\binom{n-y}{n-x}} && 0, n \neq x > y \end{aligned}$$

Note that in the statement of the theorem, $0, n$ are always points of the process.

Proof. The first step is to observe that $C(\sigma) \cup \{0, n\}$ can be obtained as a trajectory of a certain Markov chain, started at 0, with transition probabilities

$$P(i, j) = \frac{(n-j)!f(j-i)}{(n-i)!}. \quad (5.7)$$

Here we take $f(0) = 0$, so $P(i, i) = 0$ for all i . To prove (5.7), note that $C(\sigma) = \{i_1, \dots, i_k\}$ if and only if the following events E_1, \dots, E_{k+1} occur:

- E_1 : $\{\pi(1), \dots, \pi(i_1)\} = \{1, \dots, i_1\}$ and π restricted to $\{1, \dots, i_1\}$ is indecomposable.
- E_2 : $\{\pi(i_1 + 1), \dots, \pi(i_2)\} = \{i_1 + 1, \dots, i_2\}$ and π restricted to $\{i_1 + 1, \dots, i_2\}$ is indecomposable.
- ...
- E_{k+1} : $\{\pi(i_k + 1), \dots, \pi(n)\} = \{i_k + 1, \dots, n\}$ and π restricted to $\{i_k + 1, \dots, n\}$ is indecomposable.

Letting $f(n)$ denote the number of indecomposable permutations in S_n , the probability of E_1 is clearly $\frac{f(i_1)(n-i_1)!}{n!}$. The probability of E_2 given E_1 is $\frac{f(i_2-i_1)(n-i_2)!}{(n-i_1)!}$ and the probability of E_3 given E_1, E_2 is $\frac{f(i_3-i_2)(n-i_3)!}{(n-i_2)!}$, etc., as claimed.

Now [4, Thm. 1.1] implies that the point process on $C(\sigma) \cup \{0, n\}$ is determinantal with correlation kernel

$$K(x, y) = \delta_{0,x} + Q(0, x) - Q(y, x) \quad (5.8)$$

where

$$Q = P + P^2 + P^3 + \dots$$

To compute Q , let $[x^r]g(x)$ denote the coefficient of x^r in a series $g(x)$, and note that

$$\begin{aligned} P^2(i, j) &= \sum_l P(i, l)P(l, j) \\ &= \sum_l \frac{(n-l)!f(l-i)}{(n-i)!} \frac{(n-j)!f(j-l)}{(n-l)!} \\ &= \frac{(n-j)!}{(n-i)!} [x^{j-i}] \left(1 - \frac{1}{\sum_{n \geq 0} n!x^n} \right)^2. \end{aligned}$$

The last equality used (5.5). A similar computation shows that

$$P^r(i, j) = \frac{(n-j)!}{(n-i)!} [x^{j-i}] \left(1 - \frac{1}{\sum_{n \geq 0} n!x^n} \right)^r,$$

and thus

$$\begin{aligned} Q(i, j) &= \frac{(n-j)!}{(n-i)!} [x^{j-i}] \frac{\left(1 - \frac{1}{\sum_{n \geq 0} n!x^n} \right)}{\left(\sum_{n \geq 0} n!x^n \right)^{-1}} \\ &= \frac{(n-j)!}{(n-i)!} [x^{j-i}] \left(\sum_{n \geq 1} n!x^n \right). \end{aligned}$$

Thus $Q(i, j) = 1/\binom{n-i}{n-j}$ if $i < j$, and is 0 otherwise. The theorem follows from this and (5.8). \square

Remarks.

1. The connectivity set $C(\sigma)$ is a simple example of a determinantal process which is not one-dependent. Indeed, from (5.6), $P(1 \in C(\sigma)) = \frac{1!(n-1)!}{n!}$, $P(3 \in C(\sigma)) = \frac{3!(n-3)!}{n!}$, and $P(1, 3 \in C(\sigma)) = \frac{1!2!(n-3)!}{n!} \neq P(1 \in C(\sigma))P(3 \in C(\sigma))$.
2. Unlike the other point processes considered in this paper, the expected number of points tends to 0 as $n \rightarrow \infty$. Indeed, applying (5.6) gives that

$$E|C(\sigma)| = \sum_{i=1}^{n-1} P(i \in C(\sigma)) = \sum_{i=1}^{n-1} \frac{1}{\binom{n}{i}} \rightarrow 0$$

as $n \rightarrow \infty$.

Binomial posets

In Doubilet–Rota–Stanley [24], binomial posets were introduced as a unifying mechanism to “explain” the many forms of generating functions that appear in enumerative combinatorics. Briefly, \mathcal{X} is a binomial poset if for every interval $[x, y]$, the length of all maximal chains is the same (say n_{xy}) and the number of maximal chains in an interval of length n does not depend on the particular interval. This number $B(n)$ is called the factorial function. For the usual Boolean algebra of subsets, $B(n) = n!$. For the

subspaces of a vector space over a finite field, $B(n) = n!_q$ (q -factorial). There are many further examples.

Stanley [66] has shown that many enumerative formulae generalize neatly to the setting of binomial posets. Some of these developments give new determinantal point processes. Here is an example.

Example 15. (union of descent sets)

Theorem 5.5. *Pick $\sigma_1, \sigma_2, \dots, \sigma_n$ independently from Mallows model through Kendall's tau (5.3) on the symmetric group S_n . Let $S = \{s_1 < \dots < s_k\}$ be the union of their descent sets. Then*

$$P_n(S) = \det \left[\frac{1}{((s_{j+1} - s_i)!_q)^r} \right],$$

where $s_0 = 0, s_{k+1} = n$. The associated point process is stationary, one-dependent, and determinantal with $K(x, y) = k(y - x)$ for $\hat{k}(z) = \frac{1}{1 - \hat{e}(z)}$ where $\hat{e}(z) = \sum_{l=0}^{\infty} z^l / (l!_q)^r$.

Proof. The formula for $P_n(S)$ is Corollary 3.2 of [66], so the theorem follows from one of our main results proved in Section 7 (Corollary 7.6). \square

When $q = 1$ and $r = 2$, $\hat{e}(z) = I_0(2\sqrt{z})$, the classical modified Bessel function. Feller [30, §II.7] develops a host of connections with stochastic processes.

Remark. The intersection of two independent point processes with correlation functions $\rho^1(A), \rho^2(A)$ is a point process with correlation function $\rho^1(A)\rho^2(A)$. If both processes are determinantal, this is a product of determinants, but in general the intersection or union of determinantal processes is not determinantal. In Subsection 7.1, we show that all one-dependent processes are determinantal. Since the intersection of descent sets of independent permutations is one-dependent, and taking complements preserves one-dependence, this gives another proof that the union process in Theorem 5.5 is determinantal.

We have not pursued other examples but again believe there is much else to be discovered.

6 More general carries

Consider the quaternions $Q_8 = \{\pm 1, \pm i, \pm j, \pm k\}$. The center Z of Q_8 is $\{\pm 1\}$. Choose coset representatives $X = \{1, i, j, k\}$ for Z in Q_8 so any element can be uniquely represented as $g = zx$. The coset representatives are multiplied by the familiar rule $k \curvearrowright i \curvearrowright j \curvearrowright k$ so $ij = k$ and $kj = -i$, etc. If we multiply $g_1 g_2 \dots g_k = (z_1 x_1)(z_2 x_2) \dots (z_k x_k)$, the $z_i \in \{\pm 1\}$ can all be moved to the left and we must multiply $x_1 x_2 \dots x_k$, keeping track of the ‘‘carries’’, here ± 1 . Evidently, if $\{g_i\}$ are chosen uniformly at random in Q_8 , both $\{z_i\}$ and $\{x_i\}$ are independent and uniform in Z and X . Thus we have the following problem: choose X_1, X_2, \dots, X_k uniformly in X and multiply as $X_1, X_1 X_2, (X_1 X_2) X_3, \dots,$

$$\begin{aligned} &X_1 \\ &X_2 \\ &X_3 \\ &\vdots \\ &X_k \end{aligned}$$

This gives a process of remainders and carries as in Section 1.

Example 16.

$$\begin{array}{r}
 k \cdot k \\
 k \quad 1 \\
 i \cdot i \\
 i \quad 1 \\
 k \cdot k \\
 j \cdot i \\
 k \cdot j \\
 i \quad k
 \end{array}
 \quad \text{gives } (-1)^5 k = -k.$$

It is almost obvious that the carries form a stationary, one-dependent, two-block process with $P(i-1 \text{ ones in a row}) = \frac{6}{4^i}$, $2 \leq i < \infty$. Further, the remainders in the second column are independent and uniform on $\{1, i, j, k\}$ and there is a simple “descent” rule which determines the joint law of the dots (Example 18 below).

We also mention, by comparison with Example 10 in Section 4, that the carries process for the quaternions is the same point process that arises from the coordinate ring of 6 generic points in projective space \mathbb{P}^3 .

One natural generalization where all goes through is to consider a finite group G and a normal subgroup N contained in the center of the group. The factor group $F = G/N$ has elements labeled $1, \sigma, \tau, \dots$. We may choose coset representatives $t(1) = 1, t(\sigma), t(\tau), \dots$ and any $g = nt(\sigma)$, uniquely. While sometimes $t(\sigma)t(\tau) = t(\sigma\tau)$, in many cases this fails; but we may choose correction factors $f(\sigma, \tau)$ in N (often called a “factor set”) so that $t(\sigma)t(\tau) = t(\sigma\tau)f(\sigma, \tau)$. Once $t(\sigma)$ are chosen, the $f(\sigma, \tau)$ are forced.

Example 17. If $G = C_{100}$, $N = C_{10}$ (thought of as a subgroup $\{0, 10, 20, \dots, 90\}$), the natural choice of coset representatives for $G/N \cong \{0, 1, \dots, 9\}$ is $t(i) = i \in G$. Of course,

$$t(i) + t(j) = t(i + j) + f(i, j), \quad \text{with } f(i, j) = \begin{cases} 1 & \text{if } i + j \geq 10 \\ 0 & \text{if } 0 \leq i + j < 10. \end{cases}$$

It is natural to ask if the choice of cosets matters. To see that it can, consider C_{10} in C_{100} and choose coset representatives as $0, 11, 22, 43, 44, 45, 46, 47, 48, 49$. The sum of 11 and 22 requires a carry of 90.

A lovely exposition of carries as cocycles is in [42].

If $g = nt(\sigma)$ is chosen uniformly at random, then n and $t(\sigma)$ are independent and uniform. Multiplying a sequence of $g_i = n_i t_i$ can be done by first multiplying the t_i , keeping track of the carries, and then multiplying the n_i and carries in any order. Of course, here the carries are in N , not necessarily binary.

Given t_1, t_2, \dots, t_k , we may form a two-column array with t_i in the first column, the successive remainders r_1, r_2, \dots, r_k in the second column and “carries” f_1, f_2, \dots, f_{k-1} (elements in N) placed in between.

Lemma 6.1. *If coset representatives t_i , $1 \leq i \leq k$, are chosen uniformly, then*

1. *the remainder process r_i , $1 \leq i \leq k$, is uniform and independent, and*

2. the carries f_i , $1 \leq i \leq k-1$, form a stationary, one-dependent, two-block factor.

Proof.

1. Since $r_1 = t_1$, r_1 is uniform. Successive r_i are formed by multiplying r_{i-1} by t_i . There is a unique choice of t_i giving r_i , so r_i is uniform and independent of r_1, \dots, r_{i-1} .
2. Consider two successive remainders and the unique t giving rise to them:

$$\begin{array}{cc} & r_{i-1} \\ t_i & r_i \end{array}$$

Since t_i is uniquely determined, $r_{i-1}t_i = r_i f(r_{i-1}, t_i)$ is uniquely determined. It follows that the f_i process is a two-block process: generate r_1, r_2, \dots uniformly and independently, set $f_i = h(r_i, r_{i+1})$, with $h(r, r') = f(r, t)$ where $r^{-1}r' \in tN$ determines t uniquely. Because two-block processes are one-dependent, this completes the proof. \square

Corollary 6.2. *With the notation of the lemma, define a binary process B_1, \dots, B_{k-1} as*

$$B_i = \begin{cases} 1 & \text{if } f_i \neq \text{id} \\ 0 & \text{if } f_i = \text{id}. \end{cases}$$

Then $\{B_i\}$ is a stationary, one-dependent, two-block factor.

Example 18. As in Example 16, let $G = Q_8$, $N = \{\pm 1\}$, $F = G/N \cong C_2 \times C_2$. The natural choice of coset representatives $\{1, i, j, k\}$ gives rise to the following two-block representation: choose U_1, U_2, \dots uniformly and independently in $\{1, i, j, k\}$. Let $B_i = h(U_i, U_{i+1})$ with $h(1, x) = 1$ (all x), $h(x, 1) = -1$ ($x \neq 1$), $h(i, j) = h(j, k) = h(k, i) = -1$, $h(i, k) = h(k, j) = h(j, i) = 1$, $h(x, x) = 1$, ($x \neq 1$).

Example 19. (dihedral group) Let $D_8 = \langle x, y : x^2 = y^2 = (xy)^4 = 1 \rangle$. If $z = xy$, this 8-element group has center $N = \{1, z^2 = -1\}$. Choosing coset representatives $1, x, y, z$, the cosets multiply as

	1	x	y	z
1	1	x	y	z
x	x	1	z	y
y	y	$-z$	1	$-x$
z	z	$-y$	x	-1

From this, elementary manipulations show that $P(i \text{ ones in a row}) = 1/4^i$. Thus, the carries process is independent with $P(B_i = 1) = 1/4$, $P(B_i = 0) = 3/4$ for all i .

We mention that D_8 can also be represented as the extension of the normal subgroup C_4 by the factor group C_2 . Here, coset representatives for C_4 can be chosen so that there are no carries (i.e. the extension ‘‘splits’’).

Example 20. (extensions of C_2 by C_m) A central extension of $N = C_2$ by C_m is abelian. It follows that when m is odd, $G = C_{2m}$ is the only central extension of $N = C_2$ by C_m , and when m is even, there are two central extensions $C_2 \times C_m$ and C_{2m} . The extension

$C_2 \times C_m$ splits and choosing $\{(0, i), 0 \leq i \leq m-1\}$ as the coset representatives, there are no carries. For C_{2m} , with $C_2 \cong \{0, m\}$, choose coset representatives $0, 1, 2, \dots, m-1$. Thus

$$f(i, j) = \begin{cases} 1 & \text{if } i + j \geq m \\ 0 & \text{if } 0 \leq i + j < m. \end{cases}$$

As for usual addition, there is a carry if and only if there is a descent in the remainder column. Thus

$$P(B_1 = B_2 = \dots = B_{i-1} = 1) = \frac{\binom{m}{i}}{m^i}.$$

Example 21. Let $G = C_2 \times C_2 \times C_2$ and let $N = \{(0, 0, 0), (1, 1, 1)\}$. With coset representatives $(0, 0, 0), (1, 0, 0), (0, 1, 0), (1, 1, 0)$, there are never carries, but with coset representatives $(0, 0, 0), (1, 0, 0), (0, 1, 0), (0, 0, 1)$, there are carries. This example shows that the one-dependent process B_1, B_2, \dots depends not only on G and N , but also on the choice of coset representatives.

We have not embarked on a systematic study of carries for finite groups and believe that there is much more to do. We do note that by a result in the next section (Theorem 7.1), the above processes, being one-dependent, are determinantal.

The basic carries argument works for infinite groups as well. Let G be a locally compact group and H a closed normal subgroup. Suppose that H is in the center of G and that G/H is compact. Choose coset representatives $t(\sigma) \in G$ for $\sigma \in G/H$. As in the finite case, these define factor sets $f(\sigma, \tau)$ by $t(\sigma)t(\tau) = t(\sigma\tau)f(\sigma, \tau)$. Write $g = nt$. Since G/H is compact, it has an invariant probability measure. Choosing t_1, t_2, \dots independently from this measure and multiplying as above gives remainders and a carries process. Just as above, the remainders are independent and uniformly distributed in G/H and the carries process (with values in N) is a one-dependent, two-block factor process.

Example 22. A lovely instance of this set-up explains a classical identity. We begin with the motivation and then translate. Let σ be a permutation with number of descents $d(\sigma)$. It is known that, for σ chosen from the uniform distribution on S_n ,

$$P(d(\sigma) = j) = P(j \leq U_1 + \dots + U_n < j + 1). \quad (6.1)$$

On the right, U_1, U_2, \dots, U_n are independent uniforms on $[0, 1]$. The density and distribution function for $U_1 + \dots + U_n$ was derived by Laplace; see [30]. Foata [32] proved (6.1) by combining Laplace's calculation with an identity of Worpitzky. Richard Stanley [69] gives a bijective proof involving an elegant dissection of the n -dimensional hypercube. Jim Pitman [57] gives the following "proof from the book" which is a continuous version of our carries argument from Section 1: form two columns

$$\begin{array}{cc} U_1 & V_1 \\ U_2 & V_2 \\ & \vdots \\ & \vdots \\ U_n & V_n \end{array}$$

On the left are independent uniforms on $[0, 1]$. On the right are their remainders when added mod 1; so $V_1 = U_1$, $V_2 = U_1 + U_2 \pmod{1}$, \dots . The V_i are similarly independent

uniforms on $[0, 1]$. Place a dot at position i every time the partial sum $U_1 + \cdots + U_{i+1}$ crosses an integer. Call these dots carries. As in the discrete case, there is a dot at position i if and only if there is a descent $V_{i+1} < V_i$. The number of dots is the integer part of $U_1 + \cdots + U_n$ and also the number of descents, proving (6.1). Of course, the distribution of the descent *process* is the same as the carries process.

In the language of group theory, let $G = \mathbb{R}$, $N = \mathbb{Z}$, and $G/N \cong S_1$, the circle group. Choose coset representatives as $[0, 1)$ and factor sets in $\{0, 1\}$.

7 Proofs and generalizations

This section proves two of our main results for general one-dependent processes (we do not assume stationarity in this section). In Subsection 7.1, it is shown that all one-dependent point processes on \mathbb{Z} are determinantal, a result which is new even in the stationary case. Subsection 7.2 proves that a point process P on a finite set \mathcal{X} , with P given as a certain-shaped determinant, is one-dependent and determinantal. This covers quite a few examples from previous sections and is particularly useful in situations (such as Example 12) where the one-dependence is not a priori obvious.

7.1 One dependent processes are determinantal

For a random point process on a discrete set \mathcal{X} , we define the correlation function ρ by

$$\rho(A) = P\{S : S \supseteq A\}.$$

Then one dependence on (a segment of) \mathbb{Z} is equivalent to the condition that $\rho(X \cup Y) = \rho(X)\rho(Y)$ whenever $\text{dist}(X, Y) \geq 2$.

Theorem 7.1. *Any one-dependent point process on (a segment of \mathbb{Z}) is determinantal. Its correlation kernel can be written in the form $K(x, y) =$*

$$\begin{array}{ll} 0 & \text{if } x - y \geq 2 \\ -1 & \text{if } x - y = 1 \\ \sum_{r=1}^{y-x+1} (-1)^{r-1} \sum_{x=l_0 < l_1 < \cdots < l_r=y+1} \rho([l_0, l_1]) \rho([l_1, l_2]) \cdots \rho([l_{r-1}, l_r]) & \text{if } x \leq y \end{array}$$

Here the notation $[a, b)$ stands for $\{a, a+1, \dots, b-1\}$.

For example, $K(x, x) = \rho(\{x\})$, $K(x, x+1) = \rho(\{x, x+1\}) - \rho(\{x\})\rho(\{x+1\})$, etc.

Proof. By one-dependence, it is enough to verify that with K as above,

$$\det[K(x+i, x+j)]_{i,j=0}^{y-x} = \rho([x, y+1))$$

for any $x \leq y$.

We use induction on $y-x$. For $y=x$ the statement is trivial. Otherwise, one has that

$$\det[K(x+i, x+j)]_0^{y-x} = \det \begin{bmatrix} K(x, x) & K(x, x+1) & \cdots & K(x, y) \\ -1 & K(x+1, x+1) & \cdots & \vdots \\ & -1 & \ddots & \vdots \\ & & & -1 & K(y, y) \end{bmatrix}.$$

When expanding this determinant, various numbers of -1 's from the subdiagonal can be used. Observe that if we do not use the -1 in position $(i, i - 1)$, then we can compute the corresponding contribution, because if we replace that -1 by 0 , the determinant splits into the product of two each of which is computable by the induction hypothesis.

Similarly, if we insist on not using several -1 's, then the contribution is the product of several determinants. Thus we obtain by inclusion-exclusion that

$$\begin{aligned} \det[K(x+i, x+j)]_0^{y-x} &= K(x, y) + \sum_{x < l < y+1} \rho([x, l]) \rho([l, y+1]) \\ &\quad - \sum_{x < l_1 < l_2 < y+1} \rho([x, l_1]) \rho([l_1, l_2]) \rho([l_2, y+1]) + \dots \end{aligned} \quad (7.1)$$

The $K(x, y)$ term corresponds to using all -1 's, the sum over l corresponds to not using a -1 at least at location $(l, l - 1)$, the sum over l_1, l_2 corresponds to not using a -1 at least at locations $(l_j, l_j - 1), j = 1, 2$, etc. Our definition of $K(x, y)$ is such that the right hand side of (7.1) is equal to $\rho([x, y+1])$, as desired. \square

To state some corollaries of Theorem 7.1, we use the concept of particle-hole involution. Essentially, given a subset \mathcal{N} of \mathcal{X} , the involution maps a point configuration $S \subset \mathcal{X}$ to $S \triangle \mathcal{N}$ (here \triangle is the symbol for symmetric difference). This map leaves intact the particles of S outside of \mathcal{N} , and inside \mathcal{N} it loses the particles and picks up the ‘‘holes’’ (points of \mathcal{N} free of particles).

Corollary 7.2. *The class of one-dependent processes is closed under the operations of particle-hole involution on any fixed subset, intersections of independent processes, and unions of independent processes.*

Proof. For intersections, the claim follows from the definitions.

For particle-hole involutions, note that the property of being one-dependent follows from the kernel begin 0 on the second subdiagonal and below. On the other hand, for determinantal point processes the ‘‘complementation principle’’ [5, Sec. A.3] says that the particle-hole involution can be implemented by the following change in the kernel:

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix} \rightarrow \begin{bmatrix} A & B \\ -C & I - D \end{bmatrix},$$

where the block structure corresponds to the splitting into the noninverted and inverted parts. Clearly, this keeps the property of having 0 's on the second subdiagonal and below intact.

Finally, unions can be reduced to intersections by the particle-hole involution. \square

Remark. The class of determinantal point processes is *not* closed under intersections/unions.

Let us now look at the translation invariant case. Then $\rho([x, y]) = \rho_{y-x}$ is a function of $y - x$ only, and ρ_k is the chance of k consecutive ones. Set

$$R(z) = 1 + z + \sum_{k \geq 1} \rho_k z^{k+1}.$$

Corollary 7.3. *In the translation invariant case, the kernel $K(x, y) = k(y - x)$ is also translation invariant, and*

$$\sum_{n \in \mathbb{Z}} k(n)z^n = \frac{1}{1 - R(z)}.$$

Proof.

$$\begin{aligned} \frac{1}{1 - R(z)} &= -\frac{1}{z} \cdot \frac{1}{1 + \rho_1 z + \rho_2 z^2 + \dots} \\ &= -\frac{1}{z} \left[1 - \sum_{m \geq 1} \rho_m z^m + \left(\sum_{m \geq 1} \rho_m z^m \right)^2 - \dots \right] \\ &= -\frac{1}{z} + \frac{1}{z} \left[\sum_{m \geq 1} \rho_m z^m - \left(\sum_{m \geq 1} \rho_m z^m \right)^2 + \dots \right], \end{aligned}$$

which agrees with the formula of Theorem 7.1. \square

The generating function $R(z)$ also behaves well with respect to the particle-hole involution on \mathbb{Z} .

Proposition 7.4. *The particle-hole involution on \mathbb{Z} with $\mathcal{N} = \mathbb{Z}$ replaces $R(z)$ by $1/R(-z)$.*

Proof. We have

$$\frac{1}{1 - 1/R(-z)} = \frac{R(-z)}{R(-z) - 1} = 1 - \frac{1}{1 - R(-z)}.$$

Hence by Corollary 7.3, changing $R(z)$ to $\tilde{R}(z) = 1/R(-z)$ leads to the following change in the correlation kernel:

$$\tilde{k}(n) = \delta_{0,n} - (-1)^n k(n).$$

This is equivalent to $\tilde{K}(x, y) = \delta_{x,y} - \frac{(-1)^x}{(-1)^y} K(x, y)$. This is the same as $K \rightarrow I - K$, which corresponds to the particle-hole involution [5, Sec. A.3]. \square

Example 23. The descent process on \mathbb{Z} (Example 11) corresponds to $\rho_m = \frac{1}{(m+1)!}$; thus $R(z) = e^z$. The particle-hole involution is given by $\tilde{R}(z) = 1/e^{-z} = e^z$, which is the same.

Example 24. The intersection of r independent descent processes on \mathbb{Z} corresponds to $\rho_m = \frac{1}{(m+1)^r}$, hence

$$R_{\cap}^{(r)}(z) = \sum_{m \geq 0} \frac{z^m}{(m!)^r}.$$

The correlation kernel is given by $\sum_{n \in \mathbb{Z}} k(n)z^n = \frac{1}{1 - R_{\cap}^{(r)}(z)}$.

Example 25. The union of r independent descent processes is the particle-hole involution of Example 24. Thus $R_{\cup}^{(r)}(z) = 1/R_{\cap}^{(r)}(-z)$ and

$$\sum_{n \in \mathbb{Z}} k(n)z^n = \frac{1}{1 - 1/R_{\cap}^{(r)}(-z)}.$$

Replacing z by $-z$ doesn't affect correlations, and we recover Example 15.

7.2 A class of determinantal processes

For any $n = 2, 3, \dots$, consider a probability measure P_n on all subsets $S = \{s_1 < s_2 < \dots < s_k\} \subseteq [n-1]$ given by

$$P_n(S) = h(n) \det [e(s_i, s_{j+1})]_{i,j=0}^k, \quad (7.2)$$

for some $h : \mathbb{N} \rightarrow \mathbb{C}$ and $e : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{C}$ with the notation $s_0 = 0, s_{k+1} = n$. We assume that $e(i, j) = 0$ for $i > j$, and that $e(i, i) = 1, e(i, i+1) > 0$ for all i .

Theorem 7.5. *If (7.2) holds for some fixed n with $e(i, j) = 0$ for $i > j$ and $e(i, i) = 1, e(i, i+1) > 0$ for all i , then P_n is a determinantal, one-dependent process with correlation functions*

$$\rho(A) = P_n\{S : S \supseteq A\} = \det [K(a_i, a_j)]_{i,j=1}^m \quad \text{for } A = \{a_1, a_2, \dots, a_m\},$$

with correlation kernel

$$K(x, y) = \delta_{x,y} + (E^{-1})_{x,y+1}, \quad (7.3)$$

where E is the upper triangular matrix $E = [e(i-1, j)]_{i,j=1}^n$,

$$E = \begin{bmatrix} e(0,1) & e(0,2) & \dots & e(0,n) \\ & e(1,2) & \dots & e(1,n) \\ & & \ddots & \vdots \\ & & & e(n-1,n) \end{bmatrix}.$$

In addition, $h(n) = (\det E)^{-1} = (e(0,1)e(1,2)\dots e(n-1,n))^{-1}$.

Corollary 7.6. *Assume further that $e(i, j) = e(j-i)$. Then the point process is stationary. If $\hat{e}(z) = \sum_{l=0}^{\infty} e(l)z^l$, then $K(x, y) = k(y-x)$ and*

$$\hat{k}(z) := \sum_{l=-\infty}^{\infty} k(l)z^l = \frac{1}{1 - 1/\hat{e}(z)}.$$

Proof of Theorem 7.5. Set $L = [e(i-1, j) + \delta_{i-1,j}]_{i,j=1}^n$. Thus L appears as

$$L = \begin{bmatrix} e(0,1) & e(0,2) & \dots & e(0,n) \\ 1 & e(1,2) & \dots & \vdots \\ & 1 & \ddots & \vdots \\ & & 1 & e(n-1,n) \end{bmatrix}.$$

For any function $f : \{1, \dots, n-1\} \rightarrow \mathbb{C}$, by (7.2),

$$\begin{aligned}
& E \left(\prod_{s_i \in S} f(s_i) \right) \\
&= \sum_{\substack{0 < s_1 < \dots < s_k < n \\ k=0, \dots, n-1}} P_n(\{s_1 < s_2 < \dots < s_k\}) f(s_1) \dots f(s_k) \\
&= h(n) \sum_{\substack{0 < s_1 < \dots < s_k < n \\ k=0, \dots, n-1}} \det \left[L \begin{pmatrix} 1, s_1+1, \dots, s_k+1 \\ s_1, \dots, s_k, n \end{pmatrix} \right] f(s_1) \dots f(s_k) \\
&= h(n) \cdot \det \begin{bmatrix} f(1)e(0,1) & f(2)e(0,2) & \dots & f(n-1)e(0,n-1) & e(0,n) \\ f(1)-1 & f(2)e(1,2) & \dots & f(n-1)e(1,n-1) & e(1,n) \\ & f(2)-1 & \ddots & f(n-1)e(2,n-1) & e(2,n) \\ & & \ddots & \dots & \dots \\ & & & f(n-1)-1 & e(n-1,n) \end{bmatrix} \\
&= h(n) \cdot \det \left[\begin{bmatrix} 0 & & & & \\ -1 & \ddots & & & \\ & \ddots & \ddots & & \\ & & \ddots & \ddots & \\ & & & -1 & 0 \end{bmatrix} + L \begin{bmatrix} f(1) & & & & \\ & \ddots & & & \\ & & \ddots & & \\ & & & f(n-1) & \\ & & & & 1 \end{bmatrix} \right].
\end{aligned}$$

This is the generating functional of P_n . In the second equality, the determinant is of the minor of L with rows $1, s_1+1, \dots, s_k+1$ and columns s_1, \dots, s_k, n . In the third inequality, the 2^{n-1} possible summands correspond to choosing which of the first $n-1$ matrix columns use the -1 coming from $f(i)-1$ in the determinant expansion.

If $f = 1 + g$, the generating functional can be expressed in terms of the correlation functions:

$$\begin{aligned}
& 1 + \sum_{\substack{s_1 < \dots < s_m \\ m=1,2,\dots}} \rho_m(s_1, \dots, s_m) g(s_1) \dots g(s_m) \\
&= E \left[\prod_{s_i \in S} (1 + g(s_i)) \right] \\
&= h(n) \det \left(E + L \begin{bmatrix} g(1) & & & \\ & \ddots & & \\ & & g(n-1) & \\ & & & 0 \end{bmatrix} \right) \\
&= h(n) \det(E) \cdot \det \left(I + E^{-1}L \cdot \begin{bmatrix} g(1) & & & \\ & \ddots & & \\ & & g(n-1) & \\ & & & 0 \end{bmatrix} \right).
\end{aligned}$$

This holds for all g . First take $g = 0$ to see $h(n) \det(E) = 1$. Next note the expansion: if M is $n \times n$, then $\det[I + M] = \sum_S \det(M(S))$ with the sum over all 2^n subsets of

$[n]$, and $M(S)$ the minor with rows and columns in S . This gives that $\rho_m(s_1, \dots, s_m) = \det[K(s_i, s_j)]_{i,j=1}^m$, where $K(x, y) = (E^{-1}L)_{xy}$. Then

$$E^{-1}L = E^{-1} \left(\begin{bmatrix} 0 & & & & \\ 1 & \ddots & & & \\ & \ddots & \ddots & & \\ & & \ddots & \ddots & \\ & & & 1 & 0 \end{bmatrix} + E \right) = I + E^{-1} \begin{bmatrix} 0 & & & & \\ 1 & \ddots & & & \\ & \ddots & \ddots & & \\ & & \ddots & \ddots & \\ & & & 1 & 0 \end{bmatrix}, \quad \square$$

and the proof of (7.3) is complete.

Finally, we note that since $K(x, y)$ vanishes below the first subdiagonal, P_n is one-dependent.

Proof of Corollary. Here E is a Toeplitz matrix with symbol $(\hat{e}(z) - 1)/z$. It is triangular, so E^{-1} is a Toeplitz matrix with symbol $z/(\hat{e}(z) - 1)$. Thus K is a Toeplitz matrix with symbol $\hat{k}(z) = 1 + \frac{1}{\hat{e}(z)-1} = \frac{1}{1-(\hat{e}(z))^{-1}}$. \square

We also note that in the translation invariant case, there is an expression for $P_n(S)$ in terms of skew Schur functions of ribbon type.

Corollary 7.7. *As in Corollary 7.6, assume further that $e(i, j) = e(j - i)$. Let λ and μ be the partitions defined by*

$$\lambda_i = n - s_{i-1} - k + i - 1, \quad \mu_i = n - s_i - k + i - 1, \quad 1 \leq i \leq k + 1.$$

Let λ', μ' denote the transpose partitions of λ and μ and let $s_{\lambda'/\mu'}$ denote the corresponding skew-Schur function, obtained by specializing the elementary symmetric functions e_i to equal $e(i)$. Then $P_n(S) = \frac{1}{e(1)^n} s_{\lambda'/\mu'}$.

Proof. By assumption, formula (7.2) becomes

$$P_n(S) = h(n) \cdot \det(e_{s_{j+1}-s_i})_{i,j=0}^k.$$

Now use the argument of Theorem 4.2, together with the identification of the normalizing constant $h(n) = 1/e(1)^n$ in Theorem 7.5. \square

Remarks.

1. Another approach to Theorem 7.5 is via the theory of conditional L -ensembles in Borodin–Rains [6, Prop. 1.2].
2. In the translation invariant case $e(i, j) = e(j - i)$, Stanley [67, p. 90, Ex. 14] shows that

$$P_n([n - 1]) = \text{chance that all sites are occupied}$$

is the coefficient of z^n in the power series $h(n)\hat{e}(-z)^{-1}$. To prove this using symmetric function theory, note from Corollary 7.7 that

$$P_n([n - 1]) = h(n) \cdot s_{(n)}.$$

Here the Schur function $s_{(n)} = h_n$ (where h_n is the n th complete homogeneous symmetric function and $h(n)$ is the normalizing constant in (7.2)). If $E(z) = \sum_{r=0}^{\infty} e_r z^r$ is the generating function for elementary symmetric functions and $H(z) = \sum_{r=0}^{\infty} h_r z^r$, Macdonald [51, p. 21, (2.6)] shows that $H(z)E(-z) = 1$. This gives Stanley's formula since in Corollary 7.7, the value of e_r is $e(r)$.

Example 26. (Descents in a random sequence) From Fact 5 of Section 2, we are in the Toeplitz case with $e(j) = \binom{j+b-1}{b-1}$, $\hat{e}(z) = (1-z)^{-b}$. Applying Corollary 7.6 yields Theorem 3.1.

Example 27. (Descents in a uniform permutation) From MacMahon's formula (5.1), we are again in the Toeplitz case with $e(j) = 1/j!$, $\hat{e}(z) = e^z$. As in Example 3, one can replace z by $-z$ without changing determinants or the correlations functions. This proves Theorem 5.1.

Example 28. (Descents in a non-uniform permutation) From Stanley's formula (part a of Proposition 5.2), we are again in the Toeplitz case with $e(j) = 1/j!_q$ (q -factorial). An identity of Euler allows one to write

$$\hat{e}(z) = \prod_{m \geq 0} \frac{1}{1 - z(1-q)q^m}$$

when $0 < q < 1, |z| < 1$, but the elementary description of the correlation functions given in Proposition 5.2 seems more useful.

Acknowledgments

Borodin was partially supported by NSF grant DMS 0707163. Diaconis was partially supported by NSF grant DMS 0505673. Fulman was partially supported by NSF grant DMS 0802082 and NSA grant H98230-08-1-0133.

References

- [1] Ando, T., Totally positive matrices, *Linear Algebra Appl.* **90** (1987), 165–219.
- [2] Arnold, B. C., Balakrishnan, N. and Nagaraja, H. N. *A first course in order statistics*, Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics. John Wiley & Sons, New York, 1992.
- [3] Billera, L. J., Thomas, H. and van Willigenburg, S., Decomposable compositions, symmetric quasisymmetric functions and equality of ribbon Schur functions, *Adv. Math.* **204** (2006), 204-240.
- [4] Borodin, A., Loop-free Markov chains as determinantal point processes, *Ann. Inst. Henri Poincaré Probab. Stat.* **44** (2008), no. 1, 19-28.
- [5] Borodin, A., Okounkov, A. and Olshanski, G., Asymptotics of Plancherel measures for symmetric groups, *J. Amer. Math. Soc.* **13** (2000), 481-515 (electronic).
- [6] Borodin, A. and Rains, E. M., Eynard-Mehta theorem, Schur process, and their Pfaffian analogs, *J. Stat. Phys.* **121** (2005), 291-317.
- [7] Bradley, R. C., On a stationary, triple-wise independent, absolutely regular counterexample to the central limit theorem, *Rocky Mountain J. Math.* **37** (2007), 25-44.
- [8] Brenti, F., Unimodal, log-concave and Pólya frequency sequences in combinatorics, *Mem. Amer. Math. Soc.* **81** (1989), no. 413, viii+106.

- [9] Broman, E. I., One-dependent trigonometric determinantal processes are two-block-factors, *Ann. Probab.* **33** (2005), 601-609.
- [10] Burton, R. and Pemantle, R., Local characteristics, entropy and limit theorems for spanning trees and domino tilings via transfer-impedances, *Ann. Probab.* **21** (1993), 1329-1371.
- [11] Chebikin, D., Polytopes, generating functions, and new statistics related to descents and inversions in permutations, Ph.D. thesis, Department of Mathematics, MIT, 2008.
- [12] Chen, L. H. Y. and Shao, Q., Normal approximation under local dependence, *Ann. Probab.* **32** (2004), 1985-2028.
- [13] Cifarelli, D. M. and Fortini, S., A short note on one-dependent trigonometric determinantal probability measures. Technical report, Istituto di Metodi Quantitativi, Università Bocconi, 2005.
- [14] Comtet, L., *Advanced combinatorics*, D. Reidel, Dordrecht, 1974.
- [15] Critchlow, D. E., *Metric methods for analyzing partially ranked data*, Lecture Notes in Statistics, 34. Springer-Verlag, Berlin, 1985.
- [16] Cutler, A. and McShane, R., *The Trachtenberg speed system of basic mathematics*, Doubleday & Co., Garden City, N.Y., 1960.
- [17] Daley, D. J. and Vere-Jones, D., *An introduction to the theory of point processes*, Springer Series in Statistics, Springer-Verlag, New York, 1988.
- [18] Dembo, A. and Rinott, Y., Some examples of normal approximations by Stein's method, in: *Random discrete structures (Minneapolis, MN, 1993)*, pp. 25-44, IMA Vol. Math. Appl. 76, Springer, New York, 1996.
- [19] Diaconis, P., *Group representations in probability and statistics*, Institute of Mathematical Statistics Lecture Notes—Monograph Series, 11. Institute of Mathematical Statistics, Hayward, CA, 1988.
- [20] Diaconis, P. and Fulman, J., Carries, shuffling, and symmetric functions, (2008), to appear in *Adv. Appl. Math*, available at <http://www-rcf.usc.edu/~fulman/carriesymmetricAAM.pdf>.
- [21] Diaconis, P. and Hanlon, P., Eigen-analysis for some examples of the Metropolis algorithm, in: *Hypergeometric functions on domains of positivity, Jack polynomials, and applications*, Contemp. Math. 138, Amer. Math. Soc., Providence, RI, 1992, 99-117.
- [22] Diaconis, P., McGrath, M., and Pitman, J., Riffle shuffles, cycles and descents, *Combinatorica* **15** (1995), 11-29.
- [23] Diaconis, P. and Ram, A., Analysis of systematic scan Metropolis algorithms using Iwahori-Hecke algebra techniques, *Michigan Math. J.* **48** (2000), 157-190.

- [24] Doubilet, P., Rota G.-C., and Stanley, R., On the foundations of combinatorial theory. VI. The idea of generating function, in: *Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability Vol. II: Probability theory*, pp. 267-318. Univ. California Press, Berkeley, Calif., 1972.
- [25] Dyson, F. J., Statistical theory of the energy levels of complex systems. I, *J. Mathematical Phys.* **3** (1962), 140-156.
- [26] Dyson, F. J., Statistical theory of the energy levels of complex systems. II, *J. Mathematical Phys.* **3** (1962), 157-165.
- [27] Dyson, F. J., Statistical theory of the energy levels of complex systems. III, *J. Mathematical Phys.* **3** (1962), 166-175.
- [28] Edrei, A., Proof of a conjecture of Schoenberg on the generating function of a totally positive sequence, *Canadian J. Math.* **5**, (1953), 86–94.
- [29] Evans, D. L., Leemis, L. M. and Drew, J. H., The distribution of order statistics for discrete random variables with applications to bootstrapping, *INFORMS J. Comput.* **18** (2006), 19-30.
- [30] Feller, W., *An introduction to probability theory and its applications. Vol. II.*, 2nd ed., John Wiley and Sons, New York-London-Sydney, 1971.
- [31] Fligner, M. A. and Verducci, J. S. (eds.), *Probability models and statistical analyses for ranking data*, Lecture Notes in Statistics 80. Springer-Verlag, New York, 1993.
- [32] Foata, D., Distributions Eulériennes et Mahoniennes sur le groupe des permutations, in: *Higher Combinatorics*, M. Aigner, editor, page 47. Reidel, Dordrecht/Boston, 1977.
- [33] Foulkes, H. O., Enumeration of permutations with prescribed up-down and inversion sequences, *Discrete Math.* **15** (1976), 235-252.
- [34] Fröberg, R., Koszul algebras, in: *Advances in commutative ring theory (Fez, 1997)*, 337-350, Lecture Notes in Pure and Appl. Math., 205, Dekker, New York, 1999.
- [35] Fulman, J., Applications of symmetric functions to cycle and increasing subsequence structure after shuffles, *J. Algebr. Comb.* **16** (2002), 165-194.
- [36] Garsia, A. M. and Reutenauer, C., A decomposition of Solomon’s descent algebra, *Adv. Math.* **77** (1989), 189-262.
- [37] Gessel, I. and Reutenauer, C., Counting permutations with given cycle structure and descent set, *J. Combin. Theory Ser. A* **64** (1993), 189-215.
- [38] Gessel, I. and Viennot, G., Binomial determinants, paths, and hook length formulae, *Adv. in Math.* **58** (1985), 300-321.
- [39] Harper, L. H., Stirling behavior is asymptotically normal, *Ann. Math. Statist.* **38** (1967), 410-414.
- [40] Hoeffding, W. and Robbins, H., The central limit theorem for dependent random variables, *Duke Math. J.* **15** (1948), 773-780.

- [41] Hough, J. B., Krishnapur, M., Peres, Y., and Virág, B., Determinantal processes and independence, *Probab. Surv.* **3** (2006), 206-229 (electronic).
- [42] Isaksen, D., A cohomological viewpoint on elementary school arithmetic, *Amer. Math. Monthly* **109** (2002), 796-805.
- [43] Janson, S., Some pairwise independent sequences for which the central limit theorem fails, *Stochastics* **23** (1988), 439-448.
- [44] Johansson, K., Random matrices and determinantal processes, arXiv: math-ph/0510038 (2005).
- [45] Johansson, K. and Nordenstam, E., Eigenvalues of GUE minors, *Electron. J. Probab.* **11** (2006), 1342-1371 (electronic).
- [46] Kenyon, R., Lectures on dimers, available via <http://www.math.brown.edu/~rkenyon/papers/index.html>.
- [47] Lascoux, A. and Pragacz, P., Ribbon Schur functions, *European J. Combin.* **9** (1988), 561-574.
- [48] Lyons, R., Determinantal probability measures, *Publ. Math. Inst. Hautes Études Sci.* **98** (2003), 167-212.
- [49] Lyons, R. and Steif, J., Stationary determinantal processes: phase multiplicity, Bernoullicity, entropy, and domination, *Duke Math. J.* **120** (2003), 515-575.
- [50] Macchi, O., The coincidence approach to stochastic point processes, *Advances in Appl. Probability* **7** (1975), 83-122.
- [51] Macdonald, I. G., *Symmetric functions and Hall polynomials*, Second edition. The Clarendon Press, Oxford University Press, New York, 1995.
- [52] MacMahon, P. A., *Combinatory analysis. Vol. I, II (bound in one volume)*. Reprint of *An introduction to combinatory analysis* (1920) and *Combinatory analysis. Vol. I, II* (1915, 1916). Dover Publications, Mineola, NY, 2004.
- [53] Mallows, C. L., Non-null ranking models. I, *Biometrika* **44** (1957), 114-130.
- [54] Marden, J. I., *Analyzing and modeling rank data*, Monographs on Statistics and Applied Probability 64. Chapman & Hall, London, 1995.
- [55] Mehta, M. L., *Random matrices*, Third edition. Pure and Applied Mathematics (Amsterdam), 142. Elsevier/Academic Press, Amsterdam, 2004.
- [56] Peres, Y. and Virág, B., Zeros of the i.i.d. Gaussian power series: a conformally invariant determinantal process, *Acta Math.* **194** (2005), 1-35.
- [57] Pitman, J. W., Probabilistic bounds on the coefficients of polynomials with only real zeros, *J. Combin. Theory Ser. A* **77** (1997), 279-303.
- [58] Polishchuk, A. and Positselski, L., *Quadratic algebras*, University Lecture Series, 37. American Mathematical Society, Providence, RI, 2005.

- [59] Reiner, V., Signed permutation statistics, *European J. Combin.* **14** (1993), 553-567.
- [60] Reiner, V., Shaw, K. M., and van Willigenburg, S., Coincidences among skew Schur functions, *Adv. Math.* **216** (2007), 118-152.
- [61] Reutenauer, C., *Free Lie algebras*, London Mathematical Society Monographs. New Series, 7. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1993.
- [62] Shirai, T. and Takahashi, Y., Random point fields associated with certain Fredholm determinants, I. Fermion, Poisson and boson point processes, *J. Funct. Anal.* **205** (2003), 414-463.
- [63] Shirai, T. and Takahashi, Y., Random point fields associated with certain Fredholm determinants, II. Fermion shifts and their ergodic and Gibbs properties, *Ann. Probab.* **31** (2003), 1533-1564.
- [64] Solomon, L., A Mackey formula in the group ring of a Coxeter group, *J. Algebra* **41** (1976), 255-264.
- [65] Soshnikov, A. Determinantal random point fields, *Russian Math. Surveys* **55** (2000), 923-975.
- [66] Stanley, R. P., Binomial posets, Möbius inversion, and permutation enumeration, *J. Combinatorial Theory Ser. A* **20** (1976), 336-356.
- [67] Stanley, R. P., *Enumerative combinatorics. Vol. 1*. Corrected reprint of the 1986 original. Cambridge Studies in Advanced Mathematics, 49. Cambridge University Press, Cambridge, 1997.
- [68] Stanley, R. P., *Enumerative combinatorics. Vol. 2*, Cambridge Studies in Advanced Mathematics, 62. Cambridge University Press, Cambridge, 1999.
- [69] Stanley, R. P., Eulerian partitions of a unit hypercube, in: *Higher Combinatorics*, M. Aigner, editor, page 47. Reidel, Dordrecht/Boston, 1977.
- [70] Stanley, R. P., Generalized riffle shuffles and quasisymmetric functions, *Ann. Comb.* **5** (2001), 479-491.
- [71] Stanley, R. P., The descent set and connectivity set of a permutation, *J. Integer Seq.* **8** (2005), Article 05.3.8, 9 pp. (electronic).
- [72] Stembridge, J. R., Counterexamples to the poset conjectures of Neggers, Stanley, and Stembridge, *Trans. Amer. Math. Soc.* **359** (2007), 1115-1128 (electronic).
- [73] de Valk, V., *One-dependent processes: two-block factors and non-two-block factors*, CWI Tract, 85. Stichting Mathematisch Centrum, Centrum voor Wiskunde en Informatica, Amsterdam, 1994.
- [74] Wilf, H. S., *Algorithms and complexity*, second edition, A K Peters, Natick, MA, 2002.